



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

U11/U12 snRNP 20K siRNA (m): sc-154831

BACKGROUND

Small nuclear ribonucleoproteins, also known as snRNPs, combine with other proteins to form spliceosomes, a complex that catalyzes pre-mRNA splicing. There are two types of spliceosomes: U2 and U12. The U2-type spliceosome is found in all eukaryotes and excises U2-type introns, which account for the majority of pre-mRNA introns. The U12-type spliceosome removes U12-type introns, which comprise less than 1% of all human introns. The U12-type spliceosome is comprised of the U11 and U12 snRNPs as well as the U4/U6.U5 tri-snRNP. U11 and U12 bind as a U11/U12 di-snRNP complex, which recognizes the 5' splice site of the pre-mRNA during the first steps of U12-type spliceosome formation. U11/U12 snRNPs contain several proteins, including seven that are unique to the U11/U12snRNP: 65K, 59K, 48K, 35K, 31K, 25K and 20K. U11/U12 snRNP 20K (U11/U12 small nuclear ribonucleoprotein 20 kDa protein), also known as ZMAT5 (Zinc finger matrin-type protein 5), is a 170 amino acid nuclear protein that contains one C3H1-type zinc finger.

REFERENCES

1. Sillekens, P.T., Beijer, R.P., Habets, W.J. and van Verooij, W.J. 1989. Molecular cloning of the cDNA for the human U2 snRNA-specific A' protein. *Nucleic Acids Res.* 17: 1893-1906.
2. Crispino, J.D., Blencowe, B.J. and Sharp, P.A. 1994. Complementation by SR proteins of pre-mRNA splicing reactions depleted of U1 snRNP. *Science* 265: 1866-1869.
3. Blencowe, B.J., Issner, R., Nickerson, J.A. and Sharp, P.A. 1998. A coactivator of pre-mRNA splicing. *Genes Dev.* 12: 996-1009.
4. Price, S.R., Evans, P.R. and Nagai, K. 1998. Crystal structure of the spliceosomal U2B'-U2A' protein complex bound to a fragment of U2 small nuclear RNA. *Nature* 394: 645-650.
5. Eldridge, A.G., Li, Y., Sharp, P.A. and Blencowe, B.J. 1999. The SRm160/300 splicing coactivator is required for exon-enhancer function. *Proc. Natl. Acad. Sci. USA* 96: 6125-6130.
6. Lundin, M., Mikkelsen, B., Gudim, M. and Syed M. 2000. Gene structure of the U2 snRNP-specific A' protein gene from *Salmo salar*: alternative transcripts observed. *Mar. Biotechnol.* 2: 204-211.
7. Nagengast, A.A. and Salz, H.K. 2001. The *Drosophila* U2 snRNP protein U2A' has an essential function that is SNF/U2B' independent. *Nucleic Acids Res.* 29: 3841-3847.
8. Jurica, M.S., Licklider, L.J., Gygi, S.R., Grigorieff, N. and Moore, M.J. 2002. Purification and characterization of native spliceosomes suitable for three-dimensional structural analysis. *RNA* 8: 426-439.
9. Will, C.L., Schneider, C., Hossbach, M., Urlaub, H., Rauhut, R., Elbashir, S., Tuschl, T. and Lührmann, R. 2004. The human 18S U11/U12 snRNP contains a set of novel proteins not found in the U2-dependent spliceosome. *RNA* 10: 929-941.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Zmat5 (mouse) mapping to 11 A1.

PRODUCT

U11/U12 snRNP 20K siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see U11/U12 snRNP 20K shRNA Plasmid (m): sc-154831-SH and U11/U12 snRNP 20K shRNA (m) Lentiviral Particles: sc-154831-V as alternate gene silencing products.

For independent verification of U11/U12 snRNP 20K (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154831A and sc-154831B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

U11/U12 snRNP 20K siRNA (m) is recommended for the inhibition of U11/U12 snRNP 20K expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor U11/U12 snRNP 20K gene expression knockdown using RT-PCR Primer: U11/U12 snRNP 20K (m)-PR: sc-154831-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.