



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UCK1 siRNA (m): sc-154885

BACKGROUND

Uridine-cytidine kinases (UCK) have important roles for the phosphorylation of nucleoside analogs that may be important in chemotherapy of cancer. The UCK family consists of two members, UCK1 and UCK2, which are both expressed in many tumor cells. UCK1 (uridine-cytidine kinase 1), also known as URK1, uridine monophosphokinase 1 or cytidine monophosphokinase 1, is a 277 amino acid protein that is expressed in skeletal muscle, heart, liver and kidney. UCK1 uses ATP or GTP to catalyze the phosphorylation of uridine and cytidine to uridine monophosphate and cytidine monophosphate, respectively. Human UCK1 shares 92% amino acid identity with its mouse counterpart, suggesting a conserved role between species. UCK1 exists as two alternatively spliced isoforms which are encoded by a gene that maps to human chromosome 9.

REFERENCES

1. Kaneko, S., et al. 1998. Cloning, sequence analysis and expression of the basidiomycete *Lentinus edodes* gene UCK1, encoding UMP-CMP kinase, the homologue of *Saccharomyces cerevisiae* URA6 gene. *Gene* 211: 259-266.
2. Van Rompay, A.R., et al. 2001. Phosphorylation of uridine and cytidine nucleoside analogs by two human uridine-cytidine kinases. *Mol. Pharmacol.* 59: 1181-1186.
3. Shimamoto, Y., et al. 2002. Sensitivity of human cancer cells to the new anticancer ribo-nucleoside TAS-106 is correlated with expression of uridine-cytidine kinase 2. *Jpn. J. Cancer Res.* 93: 825-833.
4. Suzuki, N.N., et al. 2003. Crystallization and preliminary X-ray analysis of human uridine-cytidine kinase 2. *Acta Crystallogr. D Biol. Crystallogr.* 59: 1477-1478.
5. Miyazaki, Y., et al. 2004. Target genes of the developmental regulator PRIB of the mushroom *Lentinula edodes*. *Biosci. Biotechnol. Biochem.* 68: 1898-1905.
6. Murata, D., et al. 2004. A crucial role of uridine/cytidine kinase 2 in antitumor activity of 3'-ethynyl nucleosides. *Drug Metab. Dispos.* 32: 1178-1182.
7. Suzuki, N.N., et al. 2004. Structural basis for the specificity, catalysis, and regulation of human uridine-cytidine kinase. *Structure* 12: 751-764.
8. Online Mendelian Inheritance in Man, OMIM™. 2005. Johns Hopkins University, Baltimore, MD. MIM Number: 609328. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Uck1 (mouse) mapping to 2 B.

PRODUCT

UCK1 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UCK1 shRNA Plasmid (m): sc-154885-SH and UCK1 shRNA (m) Lentiviral Particles: sc-154885-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

UCK1 siRNA (m) is recommended for the inhibition of UCK1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

UCK1 (E-9): sc-373940 is recommended as a control antibody for monitoring of UCK1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor UCK1 gene expression knockdown using RT-PCR Primer: UCK1 (m)-PR: sc-154885-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.