



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UGT2B1 siRNA (m): sc-154899

BACKGROUND

UDP-glucuronosyltransferase isoenzymes (UGTs) catalyze the glucuronidation of small lipophilic molecules, which regulates the bioactivity and metabolic fate of a wide range of endo- and xenobiotics. Glucuronidation increases the polarity of lipophilic molecules and facilitates their entry into aqueous compartments and their ultimate excretion. In essence, glucuronidation provides a protective function by terminating or attenuating the biological activity of its substrates. The UGT2B family of isoenzymes are highly expressed in liver, but are also detected in several non-hepatic tissues, including skin, breast, prostate, intestine, placenta, and lung. Therefore, the UGT2B family may preferentially modulate steroid metabolism and excretion in addition to bile acids and xenobiotics. The human UGT2B genes localize as a cluster on chromosome 4q13.

REFERENCES

1. Monaghan, G., Clarke, D.J., Povey, S., See, C.G., Boxer, M. and Burchell, B. 1994. Isolation of a human YAC contig encompassing a cluster of UGT2 genes and its regional localization to chromosome 4q13. *Genomics* 23: 496-499.
2. Beaulieu, M., Levesque, E., Hum, D.W. and Belanger, A. 1996. Isolation and characterization of a novel cDNA encoding a human UDP-glucuronosyltransferase active on C19 steroids. *J. Biol. Chem.* 271: 22855-22862.
3. Belanger, A., Hum, D.W., Beaulieu, M., Levesque, E., Guillemette, C., Tchernof, A., Belanger, G., Turgeon, D. and Dubois, S. 1998. Characterization and regulation of UDP-glucuronosyltransferases in steroid target tissues. *J. Steroid Biochem. Mol. Biol.* 65: 301-310.
4. Beaulieu, M., Levesque, E., Hum, D.W. and Belanger, A. 1998. Isolation and characterization of a human orphan UDP-glucuronosyltransferase, UGT2B11. *Biochem. Biophys. Res. Commun.* 248: 44-50.
5. Levesque, E., Beaulieu, M., Hum, D.W. and Belanger, A. 1999. Characterization and substrate specificity of UGT2B4 (E458): a UDP-glucuronosyltransferase encoded by a polymorphic gene. *Pharmacogenetics* 9: 207-216.
6. Lampe, J.W., Bigler, J., Bush, A.C. and Potter, J.D. 2000. Prevalence of polymorphisms in the human UDP-glucuronosyltransferase 2B family: UGT2B4(D458E), UGT2B7(H268Y), and UGT2B15(D85Y). *Cancer Epidemiol. Biomarkers Prev.* 9: 329-333.
7. Gestl, S.A., Green, M.D., Shearer, D.A., Frauenhoffer, E., Tephly, T.R. and Weisz, J. 2002. Expression of UGT2B7, a UDP-glucuronosyltransferase implicated in the metabolism of 4-hydroxyestrone and all-*trans* retinoic acid, in normal human breast parenchyma and in invasive and *in situ* breast cancers. *Am. J. Pathol.* 160: 1467-1479.

CHROMOSOMAL LOCATION

Genetic locus: Ugt2b1 (mouse) mapping to 5 E1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

UGT2B1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UGT2B1 shRNA Plasmid (m): sc-154899-SH and UGT2B1 shRNA (m) Lentiviral Particles: sc-154899-V as alternate gene silencing products.

For independent verification of UGT2B1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-154899A, sc-154899B and sc-154899C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

UGT2B1 siRNA (m) is recommended for the inhibition of UGT2B1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor UGT2B1 gene expression knockdown using RT-PCR Primer: UGT2B1 (m)-PR: sc-154899-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.