



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ULK2 siRNA (m): sc-154914

## BACKGROUND

ULK1 and ULK2 (for UNC-51-like kinase) encode similar amino-terminal serine/threonine kinase domains, a proline/serine-rich (PS) domain and a species-conserved carboxyl-terminal domain. Both share homology with the UNC-51 kinase from *Caenorhabditis elegans* and the APG1 kinase in yeast, which are involved in axonal extension and growth, and autophagy, respectively. ULK1 maps to human chromosome 12q24.33 and is ubiquitously expressed. ULK2, also widely expressed, maps to mouse chromosome 11 B2 and is expected to have a similar molecular weight as ULK1 in human. ULK1 and ULK2 are thought to autophosphorylate the PS domain *in vitro*, and the significant homology among vertebrates suggest that ULK1 and ULK2 are involved in the regulation of fundamental biological processes.

## REFERENCES

- Ogura, K., et al. 1994. *Caenorhabditis elegans* unc-51 gene required for axonal elongation encodes a novel serine/threonine kinase. *Genes Dev.* 8: 2389-2400.
- Matsuura, A., et al. 1997. Apg1p, a novel protein kinase required for the autophagic process in *Saccharomyces cerevisiae*. *Gene* 192: 245-250.
- Kuroyanagi, H., et al. 1998. Human ULK1, a novel serine/threonine kinase related to UNC-51 kinase of *Caenorhabditis elegans*: cDNA cloning, expression, and chromosomal assignment. *Genomics* 51: 76-85.
- Yan, J., et al. 1998. Identification of mouse ULK1, a novel protein kinase structurally related to *C. elegans* UNC-51. *Biochem. Biophys. Res. Commun.* 246: 222-227.
- Yan, J., et al. 1999. Mouse ULK2, a novel member of the UNC-51-like protein kinases: unique features of functional domains. *Oncogene* 18: 5850-5859.
- Tomoda, T., et al. 1999. A mouse serine/threonine kinase homologous to *C. elegans* UNC-51 functions in parallel fiber formation of cerebellar granule neurons. *Neuron* 24: 833-846.

## CHROMOSOMAL LOCATION

Genetic locus: Ulk2 (mouse) mapping to 11 B2.

## PRODUCT

ULK2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ULK2 shRNA Plasmid (m): sc-154914-SH and ULK2 shRNA (m) Lentiviral Particles: sc-154914-V as alternate gene silencing products.

For independent verification of ULK2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3 nmol of lyophilized siRNA. These include: sc-154914A, sc-154914B and sc-154914C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ULK2 siRNA (m) is recommended for the inhibition of ULK2 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

ULK2 (2H4B2): sc-517299 is recommended as a control antibody for monitoring of ULK2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ULK2 gene expression knockdown using RT-PCR Primer: ULK2 (m)-PR: sc-154914-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.