



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

VAT1 siRNA (m): sc-155094

BACKGROUND

The storage and release of neurotransmitters in the nerve terminal is regulated by synaptic vesicles. In response to an intracellular increase in calcium levels, synaptic vesicles open and release neurotransmitters across the neuronal synapse, thereby propagating nerve impulses between neurons. VAT1 (vesicle amine transport protein 1) is a 393 amino acid integral membrane protein that is located within cholinergic synaptic vesicles. Expressed in tissues throughout the body, VAT1 belongs to the quinone oxidoreductase subfamily of zinc-containing alcohol dehydrogenase proteins and is thought to play a role in vesicular transport. Defects in the gene encoding VAT1 may be associated with endocrine disorders and tumorigenesis.

REFERENCES

1. Linal, M., et al. 1989. VAT-1: an abundant membrane protein from Torpedo cholinergic synaptic vesicles. *Neuron* 2: 1265-1273.
2. Peter, D., et al. 1993. Chromosomal localization of the human vesicular amine transporter genes. *Genomics* 18: 720-723.
3. Friedman, L.S., et al. 1994. The search for BRCA1. *Cancer Res.* 54: 6374-6382.
4. Miki, Y., et al. 1994. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 266: 66-71.
5. Smith, T.M., et al. 1996. Complete genomic sequence and analysis of 117 kb of human DNA containing the gene BRCA1. *Genome Res.* 6: 1029-1049.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604631. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Holt, M., et al. 2008. Synaptic vesicles are constitutively active fusion machines that function independently of Ca²⁺. *Curr. Biol.* 18: 715-722.

CHROMOSOMAL LOCATION

Genetic locus: Vat1 (mouse) mapping to 11 D.

PRODUCT

VAT1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see VAT1 shRNA Plasmid (m): sc-155094-SH and VAT1 shRNA (m) Lentiviral Particles: sc-155094-V as alternate gene silencing products.

For independent verification of VAT1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155094A, sc-155094B and sc-155094C.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

VAT1 siRNA (m) is recommended for the inhibition of VAT1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

VAT1 (E-7): sc-515705 is recommended as a control antibody for monitoring of VAT1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker[™] compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor VAT1 gene expression knockdown using RT-PCR Primer: VAT1 (m)-PR: sc-155094-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.