



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Vav3 siRNA (m): sc-155095

BACKGROUND

The Vav family of Rho guanine nucleotide exchange factors (GEFs) orchestrate signaling events following lymphocyte antigen receptor activation. Vav3, like Vav (also known as Vav1 or p95Vav), undergoes tyrosine phosphorylation downstream of T cell receptor cross-linkage, and subsequently interacts with two adaptor molecules, SLP76 and 3BP2. Following these events, however, the paths of Vav and Vav3 diverge; Vav affects IL-2 promoter activity, while Vav3 impacts gene transcription linked to serum response element (SRE). Further-more, Vav3 expression follows a cell cycle-dependent pattern, with transient upregulation occurring during mitosis. Enforced Vav3 expression leads to the appearance of multinucleate cells, implicating a role for Vav3 in the control of cytokinesis.

REFERENCES

- Romero, F., et al. 1996. p95vav associates with the nuclear protein Ku-70. *Mol. Cell. Biol.* 16: 37-44.
- Adam, L., et al. 2000. Interferon- α signaling promotes nucleus-to-cytoplasmic redistribution of p95Vav, and formation of a multisubunit complex involving Vav, Ku80, and Tyk2. *Biochem. Biophys. Res. Commun.* 267: 692-696.
- Fujikawa, K., et al. 2002. Vav3 is regulated during the cell cycle and effects cell division. *Proc. Natl. Acad. Sci. USA* 99: 4313-4318.
- Fujikawa, K., et al. 2003. Vav1/2/3-null mice define an essential role for Vav family proteins in lymphocyte development and activation but a differential requirement in MAPK signaling in T and B cells. *J. Exp. Med.* 198: 1595-1608.
- Zakaria, S., et al. 2004. Differential regulation of TCR-mediated gene transcription by Vav family members. *J. Exp. Med.* 199: 429-434.
- Dong, Z., et al. 2006. Vav3 oncogene is overexpressed and regulates cell growth and androgen receptor activity in human prostate cancer. *Mol. Endocrinol.* 20: 2315-2325.
- Sauzeau, V., et al. 2006. Vav3 proto-oncogene deficiency leads to sympathetic hyperactivity and cardiovascular dysfunction. *Nat. Med.* 12: 841-845.

CHROMOSOMAL LOCATION

Genetic locus: Vav3 (mouse) mapping to 3 F3.

PRODUCT

Vav3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Vav3 shRNA Plasmid (m): sc-155095-SH and Vav3 shRNA (m) Lentiviral Particles: sc-155095-V as alternate gene silencing products.

For independent verification of Vav3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155095A, sc-155095B and sc-155095C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Vav3 siRNA (m) is recommended for the inhibition of Vav3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Vav3 gene expression knockdown using RT-PCR Primer: Vav3 (m)-PR: sc-155095-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.