



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

XKR9 siRNA (m): sc-155378

BACKGROUND

Kell and XK are two covalently linked plasma membrane proteins that constitute the Kell blood group system, a group of antigens on the surface of red blood cells that are important determinants of blood type and targets for autoimmune or alloimmune diseases. XK is a 444 amino acid protein that spans the membrane ten times and carries the ubiquitous antigen Kx, which determines blood type. The XK (X-linked Kx blood group)-related gene family are homologs of XK. XKR9 (XK, Kell blood group complex subunit-related family, member 9), also known as XRG9, is a 373 amino acid multi-pass membrane protein belonging to the XK family. The gene encoding XRG9 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.

REFERENCES

1. Russo, D., Wu, X., Redman, C.M. and Lee, S. 2000. Expression of Kell blood group protein in nonerythroid tissues. *Blood* 96: 340-346.
2. Lee, S., Russo, D. and Redman, C.M. 2000. The Kell blood group system: Kell and XK membrane proteins. *Semin. Hematol.* 37: 113-121.
3. Lee, S., Russo, D. and Redman, C. 2000. Functional and structural aspects of the Kell blood group system. *Transfus. Med. Rev.* 14: 93-103.
4. Singleton, B.K., Green, C.A., Renaud, S., Fuhr, P., Poole, J. and Daniels, G.L. 2003. McLeod syndrome resulting from a novel XK mutation. *Br. J. Haematol.* 122: 682-685.
5. Feder, M. and Bujnicki, J.M. 2005. Identification of a new family of putative PD-(D/E)XK nucleases with unusual phylogenomic distribution and a new type of the active site. *BMC Genomics* 6: 21.
6. Starling, A., Schlesinger, D., Kok, F., Passos-Bueno, M.R., Vainzof, M. and Zatz, M. 2005. A family with McLeod syndrome and calpainopathy with clinically overlapping diseases. *Neurology* 65: 1832-1833.
7. Zeman, A., Daniels, G., Tilley, L., Dunn, M., Toplis, L., Bullock, T., Poole, J. and Blackwood, D. 2005. McLeod syndrome: life-long neuropsychiatric disorder due to a novel mutation of the XK gene. *Psychiatr. Genet.* 15: 291-293.
8. Pu, J.J., Redman, C.M., Visser, J.W. and Lee, S. 2005. Onset of expression of the components of the Kell blood group complex. *Transfusion* 45: 969-974.
9. Calenda, G., Peng, J., Redman, C.M., Sha, Q., Wu, X. and Lee, S. 2006. Identification of two new members, XPLAC and XTES, of the XK family. *Gene* 370: 6-16.

CHROMOSOMAL LOCATION

Genetic locus: Xkr9 (mouse) mapping to 1 A3.

PRODUCT

XKR9 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see XKR9 shRNA Plasmid (m): sc-155378-SH and XKR9 shRNA (m) Lentiviral Particles: sc-155378-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

XKR9 siRNA (m) is recommended for the inhibition of XKR9 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor XKR9 gene expression knockdown using RT-PCR Primer: XKR9 (m)-PR: sc-155378-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.