



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ZCCHC18 siRNA (m): sc-155476

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZCCHC18 (zinc finger, CCHC domain containing 18), also known as SIZN2 or PNMA7B, is a 403 amino acid protein that belongs to the ZCCHC12 family and contains one CCHC-type zinc finger. The gene encoding ZCCHC18 maps to human chromosome Xq22.2.

REFERENCES

- Ross, M.T., Grafham, D.V., Coffey, A.J., Scherer, S., McLay, K., Muzny, D., Platzer, M., Howell, G.R., Burrows, C., Bird, C.P., Frankish, A., Lovell, F.L., Howe, K.L., Ashurst, J.L., Fulton, R.S., Sudbrak, R., Wen, G., et al. 2005. The DNA sequence of the human X chromosome. *Nature* 434: 325-337.
- Cho, G., Bhat, S.S., Gao, J., Collins, J.S., Rogers, R.C., Simensen, R.J., Schwartz, C.E., Golden, J.A. and Srivastava, A.K. 2008. Evidence that SIZN1 is a candidate X-linked mental retardation gene. *Am. J. Med. Genet. A* 146A: 2644-2650.
- Cho, G., Lim, Y., Zand, D. and Golden, J.A. 2008. Szn1 is a novel protein that functions as a transcriptional coactivator of bone morphogenic protein signaling. *Mol. Cell. Biol.* 28: 1565-1572.
- Li, H., Liu, Q., Hu, X., Feng, D., Xiang, S., He, Z., Hu, X., Zhou, J., Ding, X., Zhou, C. and Zhang, J. 2009. Human ZCCHC12 activates AP-1 and CREB signaling as a transcriptional co-activator. *Acta Biochim. Biophys. Sin.* 41: 535-544.
- Cho, G., Lim, Y. and Golden, J.A. 2009. SUMO interaction motifs in Szn1 are required for promyelocytic leukemia protein nuclear body localization and for transcriptional activation. *J. Biol. Chem.* 284: 19592-19600.
- Cho, G., Lim, Y. and Golden, J.A. 2011. XLMR candidate mouse gene, Zcchc12 (Szn1) is a novel marker of Cajal-Retzius cells. *Gene Expr. Patterns* 11: 216-220.

CHROMOSOMAL LOCATION

Genetic locus: Zcchc18 (mouse) mapping to X F1.

PRODUCT

ZCCHC18 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZCCHC18 shRNA Plasmid (m): sc-155476-SH and ZCCHC18 shRNA (m) Lentiviral Particles: sc-155476-V as alternate gene silencing products.

For independent verification of ZCCHC18 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155476A, sc-155476B and sc-155476C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZCCHC18 siRNA (m) is recommended for the inhibition of ZCCHC18 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZCCHC18 gene expression knockdown using RT-PCR Primer: ZCCHC18 (m)-PR: sc-155476-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.