



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ZFP26 siRNA (m): sc-155539

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. Zinc finger protein 26 (ZFP26), also known as Zfp70 or KRAB15, is a 861 amino acid protein belonging to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family. ZNF26 contains 23 C<sub>2</sub>H<sub>2</sub>-type zinc fingers and one KRAB domain. Localized to the nucleus, ZNF26 is involved in transcriptional regulation.

## REFERENCES

1. Chowdhury, K., Rohdewohld, H. and Gruss, P. 1988. Specific and ubiquitous expression of different Zn finger protein genes in the mouse. *Nucleic Acids Res.* 16: 9995-10011.
2. Ihle, J.N., Smith-White, B., Sisson, B., Parker, D., Blair, D.G., Schultz, A., Kozak, C., Lunsford, R.D., Askew, D. and Weinstein, Y. 1989. Activation of the c-H-Ras proto-oncogene by retrovirus insertion and chromosomal rearrangement in a Moloney leukemia virus-induced T-cell leukemia. *J. Virol.* 63: 2959-2966.
3. Bellefroid, E.J., Marine, J.C., Matera, A.G., Bourguignon, C., Desai, T., Healy, K.C., Bray-Ward, P., Martial, J.A., Ihle, J.N. and Ward, D.C. 1995. Emergence of the ZNF91 Krüppel-associated box-containing zinc finger gene family in the last common ancestor of anthropoidea. *Proc. Natl. Acad. Sci. USA* 92: 10757-10761.
4. Marine, J.C., Gilbert, D.J., Bellefroid, E.J., Martial, J.A., Ihle, J.N., Copeland, N.G. and Jenkins, N.A. 1996. Chromosomal location of fifteen unique mouse KRAB-containing zinc finger loci. *Mamm. Genome* 7: 413-416.
5. Agata, Y., Matsuda, E. and Shimizu, A. 1998. Rapid and efficient cloning of cDNAs encoding Krüppel-like zinc finger proteins by degenerate PCR. *Gene* 213: 55-64.
6. Gray, P.A., Fu, H., Luo, P., Zhao, Q., Yu, J., Ferrari, A., Tenzen, T., Yuk, D.I., Tsung, E.F., Cai, Z., Alberta, J.A., Cheng, L.P., Liu, Y., Stenman, J.M., Valerius, M.T., Billings, N., Kim, H.A., Greenberg, M.E., McMahon, A.P., et al. 2004. Mouse brain organization revealed through direct genome-scale TF expression analysis. *Science* 306: 2255-2257.
7. Millien, G., Beane, J., Lenburg, M., Tsao, P.N., Lu, J., Spira, A. and Ramirez, M.I. 2008. Characterization of the mid-foregut transcriptome identifies genes regulated during lung bud induction. *Gene Expr. Patterns* 8: 124-139.
8. Fu, H., Cai, J., Clevers, H., Fast, E., Gray, S., Greenberg, R., Jain, M.K., Ma, Q., Qiu, M., Rowitch, D.H., Taylor, C.M. and Stiles, C.D. 2009. A genome-wide screen for spatially restricted expression patterns identifies transcription factors that regulate glial development. *J. Neurosci.* 29: 11399-11408.

## CHROMOSOMAL LOCATION

Genetic locus: Zfp26 (mouse) mapping to 9 A3.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PRODUCT

ZFP26 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZFP26 shRNA Plasmid (m): sc-155539-SH and ZFP26 shRNA (m) Lentiviral Particles: sc-155539-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZFP26 siRNA (m) is recommended for the inhibition of ZFP26 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZFP26 gene expression knockdown using RT-PCR Primer: ZFP26 (m)-PR: sc-155539-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.