



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ZNF300 siRNA (m): sc-155681

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF300 (zinc finger protein 300) is a 604 amino acid nuclear protein that belongs to the Krüppel C₂H₂-type zinc-finger protein family. Containing 12 C₂H₂-type zinc fingers and a KRAB domain, ZNF300 is expressed mostly in heart, skeletal muscle and brain. ZNF300 is suggested to be a regulator of the human IL-2Rβ promoter and may be involved in promoting tumor development by modulating the NFκB pathway. ZNF300 exists as three alternatively spliced isoforms and is encoded by a gene located on human chromosome 5q33.1.

REFERENCES

1. Payre, F. and Vincent, A. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. *FEBS Lett.* 234: 245-250.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Rosenfeld, R. and Margalit, H. 1993. Zinc fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs. *J. Biomol. Struct. Dyn.* 11: 557-570.
4. Laity, J.H., Lee, B.M. and Wright, P.E. 2001. Zinc finger proteins: new insights into structural and functional diversity. *Curr. Opin. Struct. Biol.* 11: 39-46.
5. Gou, D., Wang, J., Gao, L., Sun, Y., Peng, X., Huang, J. and Li, W. 2004. Identification and functional analysis of a novel human KRAB/C₂H₂ zinc finger gene ZNF300. *Biochim. Biophys. Acta* 1676: 203-209.
6. Cao, Y., Li, J.X., Ji, C.N., Xu, X.W. and Wu, M. 2007. Molecular cloning and characterization of a novel splice variant of human ZNF300 gene, which expressed highly in testis. *DNA Seq.* 18: 312-315.
7. Liu, J. and Stormo, G.D. 2008. Context-dependent DNA recognition code for C₂H₂ zinc-finger transcription factors. *Bioinformatics* 24: 1850-1857.
8. Xue, L., Qiu, H., Ma, J., Guo, M. and Li, W. 2010. ZNF300, a recently identified human transcription factor, activates the human IL-2Rβ promoter through the overlapping ZNF300/EGR1 binding site. *Cell. Mol. Biol. Lett.* 15: 530-540.
9. Wang, T., Wang, X.G., Xu, J.H., Wu, X.P., Qiu, H.L., Yi, H. and Li, W.X. 2011. Overexpression of the human ZNF300 gene enhances growth and metastasis of cancer cells through activating NF-κB pathway. *J. Cell. Mol. Med.* 16: 1134-1145.

CHROMOSOMAL LOCATION

Genetic locus: Zfp300 (mouse) mapping to X A2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF300 siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF300 shRNA Plasmid (m): sc-155681-SH and ZNF300 shRNA (m) Lentiviral Particles: sc-155681-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF300 siRNA (m) is recommended for the inhibition of ZNF300 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF300 gene expression knockdown using RT-PCR Primer: ZNF300 (m)-PR: sc-155681-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.