



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# FXR siRNA (m): sc-155894

## BACKGROUND

The steroid receptor superfamily acts through direct association with DNA sequences known as hormone response elements (HREs) and binds DNA as either homo- or heterodimers. The promiscuous mediator of heterodimerization, RXR, is the receptor for 9-*cis* retinoic acid, and dimerizes with VDR, TR, PPAR and several novel receptors, including LXR (also referred to as RLD-1) and FXR. FXR and LXR fall into a category of proteins termed "orphan receptors" because of their lack of a defined function and, in the case of LXR, the lack of a defined ligand. FXR has been shown to bind a class of lipid molecules called farnesoids. LXR/RXR heterodimers have highest affinity for DR-4 DNA elements while FXR/RXR heterodimers bind IR-1 elements. Both LXR/RXR and FXR/RXR heterodimers retain their responsiveness to 9-*cis* retinoic acid.

## REFERENCES

1. Zechel, C., Shen, X.Q., Chen, J.Y., Chen, Z.P., Chambon, P. and Gronemeyer, H. 1994. The dimerization interfaces formed between the DNA binding domains of RXR, RAR and TR determine the binding specificity and polarity of the full-length receptors to direct repeats. *EMBO J.* 13: 1425-1433.
2. Forman, B.M., Goode, E., Chen, J., Oro, A.E., Bradley, D.J., Perlmann, T., Noonan, D.J., Burka, L.T., McMorris, T., Lamph, W.W., Evans, R.M. and Weinberger, C. 1995. Identification of a nuclear receptor that is activated by farnesol metabolites. *Cell* 81: 687-693.
3. Mangelsdorf, D.J., Thummel, C., Beato, M., Herrlich, P., Schütz, G., Umesonon K., Blumberg, B., Kastner, P., Mark, M., Chambon, P. and Evans, R.M. 1995. The nuclear receptor superfamily: the second decade. *Cell* 83: 835-839.
4. Mangelsdorf, D.J. and Evans, R.M. 1995. The RXR heterodimers and orphan receptors. *Cell* 83: 841-850.
5. Willy, P.J., Umesono, K., Ong, E.S., Evans, R.M., Heyman, R.A. and Mangelsdorf, D.J. 1995. LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev.* 9: 1033-1045.
6. Leblanc, B.P. and Stunnenberg, H.G. 1995. 9-*cis* retinoic acid signaling: changing partners causes some excitement. *Genes Dev.* 9: 1811-1816.

## CHROMOSOMAL LOCATION

Genetic locus: Nr1h4 (mouse) mapping to 10 C2.

## PRODUCT

FXR siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FXR shRNA Plasmid (m): sc-155894-SH and FXR shRNA (m) Lentiviral Particles: sc-155894-V as alternate gene silencing products.

For independent verification of FXR (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155894A, sc-155894B and sc-155894C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

FXR siRNA (m) is recommended for the inhibition of FXR expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FXR gene expression knockdown using RT-PCR Primer: FXR (m)-PR: sc-155894-PR (20  $\mu$ l, 569 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Dong, R., Wang, J., Gao, X., Wang, C., Liu, K., Wu, J., Liu, Z., Sun, H., Ma, X. and Meng, Q. 2019. Yangonin protects against estrogen-induced cholestasis in a farnesoid X receptor-dependent manner. *Eur. J. Pharmacol.* 857: 172461.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.