



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

FXR β siRNA (m): sc-155895

BACKGROUND

NR (nuclear receptor) proteins comprise a large family of nuclear hormone receptor transcription factors that are characterized by discrete domains which function in DNA and ligand binding. FXR β (farnesoid X receptor β), also known as Nr1h5 (nuclear receptor subfamily 1, group H, member 5), is a 505 amino acid murine protein that belongs to the NR family. Existing as multiple alternatively spliced isoforms, FXR β is coexpressed with FXR (farnesoid X receptor) in both adult and embryonic tissues where it stimulates transcription, specifically by dimerizing with RXR α (retinoid X receptor, α) and acting as a receptor for 9-*cis*-retinoic acid. Additionally, FXR β may function as a receptor for lanosterol, an intermediate in cholesterol biosynthesis, suggesting a role for FXR β in cholesterol metabolism.

REFERENCES

- Laffitte, B.A., Kast, H.R., Nguyen, C.M., Zavacki, A.M., Moore, D.D. and Edwards, P.A. 2000. Identification of the DNA binding specificity and potential target genes for the farnesoid X-activated receptor. *J. Biol. Chem.* 275: 10638-10647.
- Redinger, R.N. 2003. The role of the enterohepatic circulation of bile salts and nuclear hormone receptors in the regulation of cholesterol homeostasis: bile salts as ligands for nuclear hormone receptors. *Can. J. Gastroenterol.* 17: 265-271.
- Kok, T., Hulzebos, C.V., Wolters, H., Havinga, R., Agellon, L.B., Stellaard, F., Shan, B., Schwarz, M. and Kuipers, F. 2003. Enterohepatic circulation of bile salts in farnesoid X receptor-deficient mice: efficient intestinal bile salt absorption in the absence of ileal bile acid-binding protein. *J. Biol. Chem.* 278: 41930-41937.
- Otte, K., Kranz, H., Kober, I., Thompson, P., Hoefler, M., Haubold, B., Rimmel, B., Voss, H., Kaiser, C., Albers, M., Cheruvallath, Z., Jackson, D., Casari, G., Koegl, M., Pääbo, S., Mous, J., Kremoser, C. and Deuschle, U. 2003. Identification of farnesoid X receptor beta as a novel mammalian nuclear receptor sensing lanosterol. *Mol. Cell. Biol.* 23: 864-872.
- Rizzo, G., Renga, B., Mencarelli, A., Pellicciari, R. and Fiorucci, S. 2005. Role of FXR in regulating bile acid homeostasis and relevance for human diseases. *Curr. Drug Targets Immune Endocr. Metabol. Disord.* 5: 289-303.
- Cai, S.Y., Xiong, L., Wray, C.G., Ballatori, N. and Boyer, J.L. 2007. The farnesoid X receptor FXR α /NR1H4 acquired ligand specificity for bile salts late in vertebrate evolution. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 293: R1400-R1409.

CHROMOSOMAL LOCATION

Genetic locus: Nr1h5 (mouse) mapping to 3 F2.2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

FXR β siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see FXR β shRNA Plasmid (m): sc-155895-SH and FXR β shRNA (m) Lentiviral Particles: sc-155895-V as alternate gene silencing products.

For independent verification of FXR β (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155895A, sc-155895B and sc-155895C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

FXR β siRNA (m) is recommended for the inhibition of FXR β expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor FXR β gene expression knockdown using RT-PCR Primer: FXR β (m)-PR: sc-155895-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.