



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# L-type Ca<sup>++</sup> CP $\gamma$ 3 siRNA (m): sc-155914

## BACKGROUND

Voltage-dependent calcium channels are important for the release of neurotransmitters into neurons. L-type (long lasting current) voltage-dependent calcium channels are composed of four subunits: an  $\alpha$ 1 subunit, a  $\beta$  subunit, a  $\gamma$  subunit and an  $\alpha$ 2/ $\delta$  subunit. The  $\gamma$  subunit is made of eight distinct proteins (designated L-type Ca<sup>++</sup> CP  $\gamma$ 1- $\gamma$ 8) and functions by influencing the properties of the calcium current. L-type Ca<sup>++</sup> CP  $\gamma$ 3, also known as CACNG3 or Cacng2, is a 315 amino acid multi-pass membrane protein that belongs to the CACNG family. As one of the eight  $\gamma$  subunits, L-type Ca<sup>++</sup> CP  $\gamma$ 3 is thought to stabilize the calcium current when the calcium channel is in a closed (inactivated) state. Defects in the gene encoding L-type Ca<sup>++</sup> CP  $\gamma$ 3 may be associated with familial infantile convulsive disorder with paroxysmal choreoathetosis, an autosomal dominant neurological disorder.

## REFERENCES

1. Powers, P.A., et al. 1993. Molecular characterization of the gene encoding the  $\gamma$  subunit of the human skeletal muscle 1,4-dihydropyridine-sensitive Ca<sup>2+</sup> channel (CACNLG), cDNA sequence, gene structure, and chromosomal location. *J. Biol. Chem.* 268: 9275-9279.
2. Burgess, D.L., et al. 1999. Identification of three novel Ca<sup>2+</sup> channel  $\gamma$  subunit genes reveals molecular diversification by tandem and chromosome duplication. *Genome Res.* 9: 1204-1213.
3. Black, J.L. and Lennon, V.A. 1999. Identification and cloning of putative human neuronal voltage-gated calcium channel  $\gamma$ -2 and  $\gamma$ -3 subunits: neurologic implications. *Mayo Clin. Proc.* 74: 357-361.
4. Burgess, D.L., et al. 2001. A cluster of three novel Ca<sup>2+</sup> channel  $\gamma$  subunit genes on chromosome 19q13.4: evolution and expression profile of the  $\gamma$  subunit gene family. *Genomics* 71: 339-350.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606403. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Moss, F.J., et al. 2003. Human neuronal stargazin-like proteins,  $\gamma$ 2,  $\gamma$ 3 and  $\gamma$ 4; an investigation of their specific localization in human brain and their influence on CaV2.1 voltage-dependent calcium channels expressed in *Xenopus* oocytes. *BMC Neurosci.* 4: 23.

## CHROMOSOMAL LOCATION

Genetic locus: Cacng3 (mouse) mapping to 7 F3.

## PRODUCT

L-type Ca<sup>++</sup> CP  $\gamma$ 3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see L-type Ca<sup>++</sup> CP  $\gamma$ 3 shRNA Plasmid (m): sc-155914-SH and L-type Ca<sup>++</sup> CP  $\gamma$ 3 shRNA (m) Lentiviral Particles: sc-155914-V as alternate gene silencing products.

For independent verification of L-type Ca<sup>++</sup> CP  $\gamma$ 3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-155914A, sc-155914B and sc-155914C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

L-type Ca<sup>++</sup> CP  $\gamma$ 3 siRNA (m) is recommended for the inhibition of L-type Ca<sup>++</sup> CP  $\gamma$ 3 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

L-type Ca<sup>++</sup> CP  $\gamma$ 3 (A-07): sc-81889 is recommended as a control antibody for monitoring of L-type Ca<sup>++</sup> CP  $\gamma$ 3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor L-type Ca<sup>++</sup> CP  $\gamma$ 3 gene expression knockdown using RT-PCR Primer: L-type Ca<sup>++</sup> CP  $\gamma$ 3 (m)-PR: sc-155914-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.