



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

OPN siRNA (bovine): sc-156029

BACKGROUND

Osteopontin (OPN), also designated bone sialoprotein 1, urinary stone protein, SPP-1, ETA-1, nephropontin and uropontin, is an extracellular matrix cell adhesion phosphoglycoprotein. OPN is deposited into unmineralized matrix prior to calcification, leading to localization at various tissue interfaces including cement lines, lamina limitans and between collagen fibrils of fully matured hard tissues. While OPN is a major product of osteoblasts, it is also synthesized by brain and kidney cells. OPN isolated from or secreted by various tissues ranges in molecular weight due to posttranslational modifications. OPN functions as a substrate for transglutaminase and is involved in cell adhesion, chemoattraction and immunomodulation.

REFERENCES

- Butler, W.T. 1989. The nature and significance of osteopontin. *Connect. Tissue Res.* 23: 123-136.
- Singh, R.P., et al. 1990. Definition of a specific interaction between the early T lymphocyte activation 1 (ETA-1) protein and murine macrophages *in vitro* and its effect upon macrophages *in vivo*. *J. Exp. Med.* 171: 1931-1942.
- Prince, C.W., et al. 1991. Osteopontin, a substrate for transglutaminase and Factor XIII activity. *Biochem. Biophys. Res. Commun.* 177: 1205-1210.
- Denhardt, D.T., et al. 1993. Osteopontin: a protein with diverse functions. *FASEB J.* 7: 1475-1482.
- Butler, W.T. 1995. Structural and functional domains of osteopontin. *Ann. N.Y. Acad. Sci.* 760: 6-11.
- Weber, G.F., et al. 1996. The immunology of Eta-1/osteopontin. *Cytokine Growth Factor Rev.* 7: 241-248.
- McKee, M.D., et al. 1996. Osteopontin at mineralized tissue interfaces in bone, teeth and osseointegrated implants: ultrastructural distribution and implications for mineralized tissue formation, turnover and repair. *Microsc. Res. Tech.* 33: 141-164.
- Nemir, M., et al. 1998. Normal rat kidney cells secrete both phosphorylated and nonphosphorylated forms of osteopontin showing different physiological properties. *J. Biol. Chem.* 264: 18202-18208.

CHROMOSOMAL LOCATION

Genetic locus: SPP1 (bovine) mapping to 6.

PRODUCT

OPN siRNA (bovine) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see OPN shRNA Plasmid (bovine): sc-156029-SH and OPN shRNA (bovine) Lentiviral Particles: sc-156029-V as alternate gene silencing products.

For independent verification of OPN (bovine) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-156029A, sc-156029B and sc-156029C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

OPN siRNA (bovine) is recommended for the inhibition of OPN expression in bovine cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor OPN gene expression knockdown using RT-PCR Primer: OPN (bovine)-PR: sc-156029-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.