



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# CB2 siRNA (r): sc-270169

## BACKGROUND

The cannabinoid receptors (CB1 and CB2) are G protein-coupled receptors that inhibit adenylate cyclase activity in response to psychoactive cannabinoids. CB1 is expressed in brain tissue and, in low levels, in testis. CB2 has been shown to be expressed only by cells of the immune system, specifically by HL-60 cells. The cannabinoid receptors mediate most of the cannabinoid-induced responses in a dose-dependent, stereoselective manner. This response system is thought to be involved in specific brain functions, such as nociception, control of movement, memory, and neuroendocrine regulation as well as having a possible role in brain development. In addition, CB1 may mediate the addictive behavior involved with the use of psychoactive cannabinoids, such as THC in marijuana.

## REFERENCES

1. Matsuda, L.A., et al. 1990. Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 346: 561-564.
2. Gerard, C.M., et al. 1991. Molecular cloning of a human cannabinoid receptor which is also expressed in testis. *Biochem. J.* 179: 129-134.
3. Munro, S., et al. 1993. Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 365: 61-65.
4. Shire, D., et al. 1996. Molecular cloning, expression and function of the murine CB2 peripheral cannabinoid receptor. *Biochim. Biophys. Acta* 1307: 132-136.
5. Ledent, C., et al. 1999. Unresponsiveness to cannabinoids and reduced addictive effects of opiates in CB1 receptor knockout mice. *Science* 283: 401-404.
6. Sugiura, T., et al. 2000. Evidence that 2-arachidonoylglycerol but not N-palmitoylethanolamine or anandamide is the physiological ligand for the cannabinoid receptor ligands in HL-60 cells. *J. Biol. Chem.* 275: 605-612.
7. Fernandez-Ruiz, J., et al. 2000. The endogenous cannabinoid system and brain development. *Trends Neurosci.* 23: 14-20.

## CHROMOSOMAL LOCATION

Genetic locus: Cnr2 (rat) mapping to 5q36.

## PRODUCT

CB2 siRNA (r) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CB2 shRNA Plasmid (r): sc-270169-SH and CB2 shRNA (r) Lentiviral Particles: sc-270169-V as alternate gene silencing products.

For independent verification of CB2 (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270169A and sc-270169B.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CB2 siRNA (r) is recommended for the inhibition of CB2 expression in rat cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CB2 gene expression knockdown using RT-PCR Primer: CB2 (r)-PR: sc-270169-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.