



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# ADAMTS-9 siRNA (r): sc-270181

## BACKGROUND

ADAMTS (a disintegrin and metalloproteinase domain, with Thrombospondin type-1 modules) is a family of zinc-dependent proteases that are implicated in a variety of normal and pathological conditions, including arthritis and cancer. ADAMTS protein family members contain an amino-terminal propeptide domain, a metalloproteinase domain, a disintegrin-like domain and a carboxy-terminus that contains a varying number of thrombospondin type-1 (TSP-1) motifs. ADAMTS genes are primarily expressed in fetal tissues, including the lung, kidney and liver. The human ADAMTS9 gene maps to chromosome 3p14.1 and encodes a deduced 1,1471 amino acid protein that is expressed in ovary, pancreas, heart, lung, placenta, adult kidney and fetal tissues. Human chromosome 3p14.1 is a region that is known to contain deletions and rearrangements in renal cell carcinomas, breast cancers, uterine cervical carcinomas and vulvar carcinomas.

## REFERENCES

1. Tang, B.L. and Hong, W. 1999. ADAMTS: a novel family of proteases with an ADAM protease domain and thrombospondin 1 repeats. *FEBS Lett.* 445: 223-225.
2. Online Mendelian Inheritance in Man, OMIM™. 2000. Johns Hopkins University, Baltimore, MD. MIM Number: 605175. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Clark, M.E., Kelner, G.S., Turbeville, L.A., Boyer, A., Arden, K.C. and Maki, R.A. 2000. ADAMTS9, a novel member of the ADAM-TS/metalloproteinase gene family. *Genomics* 67: 343-350.
4. Tang, B.L. 2001. ADAMTS: a novel family of extracellular matrix proteases. *Int. J. Biochem. Cell Biol.* 33: 33-44.

## CHROMOSOMAL LOCATION

Genetic locus: Adamts9 (rat) mapping to 4q34.

## PRODUCT

ADAMTS-9 siRNA (r) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ADAMTS-9 shRNA Plasmid (r): sc-270181-SH and ADAMTS-9 shRNA (r) Lentiviral Particles: sc-270181-V as alternate gene silencing products.

For independent verification of ADAMTS-9 (r) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270181A, sc-270181B and sc-270181C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ADAMTS-9 siRNA (r) is recommended for the inhibition of ADAMTS-9 expression in rat cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ADAMTS-9 gene expression knockdown using RT-PCR Primer: ADAMTS-9 (r)-PR: sc-270181-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.