



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Monosan® Pronase Solution

**REF** / Cat. No.: MON-APP154 250 ml

### Instructions for use

#### Intended use

Monosan® Pronase Solution is a ready-to-use solution developed for enzymatic epitope retrieval on formalin-fixed tissue sections on slides. This procedure (sometimes called PIER, *Protease Induced Epitope Retrieval*) is primarily used in immunohistochemical staining procedures.

Monosan® Pronase solution is intended for research use only, not for drug, diagnostic or other use.

#### Summary and explanations

Immunohistochemical staining procedures consist of sequential incubation steps with blocking solutions, antibodies and secondary reagents, enzymes and chromogenic substrates carried out on tissue sections. These tissue sections are mostly prepared out of formalin-fixed paraffin-embedded tissue blocks. Cellular structures are very effectively stabilised by formalin fixation which results in optimal morphological preservation of the sample.

On the other hand the formalin fixation leads to strong cross-links between proteins. This means that epitopes of antigens are being masked and often are no longer accessible for primary antibodies. In order to enable primary antibodies to bind to antigens the epitopes have to be recovered.

Enzymatic digestion with proteolytic enzymes (PIER) restores structures of the epitopes making them more accessible to specific antibodies. Heat induced epitope retrieval (HIER) in buffer solutions of different compositions and pH-values is another way of recovering epitopes. The primary antibody used determines the appropriate method.

#### Principle of the method

Monosan® Pronase Solution is a ready-to-use pronase solution in TBS buffer for enzymatic epitope retrieval.

#### Reagent provided

**REF** / Cat. No. MON-APP154  
250 ml **Monosan® Pronase Solution (Ready-To-Use)**

#### Storage and handling

The solution should be stored in aliquots at -20 °C without further dilution. Under these conditions the solution is stable up to the expiry date indicated on the label. Do not use product after the expiry date.

A positive and a negative control have to be carried out in parallel to the test material. If you observe unusual staining or other deviations from the expected results which could possibly be caused by this reagent, please contact Monosans' technical support or your local distributor.

#### Precautions

Monosan® Pronase Solution is not considered hazardous material in terms of directive 67/548/EWG. No Material safety data sheet and no marking according to the EG-directive are necessary.

#### Reagent preparation

Monosan® Pronase solution is ready-to-use and should be at room temperature prior to use.

## Procedure

Monosan® Pronase Solution is suitable for enzymatic epitope retrieval carried out after the dewaxing and rehydration of the tissue sections.

- 1) Cover deparaffinised and rehydrated tissue sections with ready-to-use Monosan® Pronase Solution.
- 2) Incubate for 10 - 10 minutes. The optimal incubation time needs to be elaborated by the operator.
- 3) Rinse carefully (3 x), first with distilled water followed by buffer.
- 4) Proceed with immunohistological staining as usual.

## Quality control

We recommend carrying out a positive and a negative control with every staining run. The positive control permits the validation of appropriate processing of the sample. If the negative control has a positive result, this points to unspecific staining. Please refer to the instructions of the detection system for guidance on general quality control procedures.

## Troubleshooting

If you observe unusual staining or other deviations from the expected results please read these instructions carefully, contact Monosans' technical support or your local distributor. Also refer to the instructions of the detection systems for guidance on general troubleshooting.

## Expected results

During the reaction of the substrate with horse radish peroxidase or alkaline phosphatase in the presence of a chromogen, a coloured precipitate is formed at the location of the bound primary antibody. This reaction only takes place if the target antigen is existent in the tissue. The chromogen used determines the colour of the precipitate. The analysis is carried out using a light microscope.

## Limitations of the procedure

Immunohistochemistry is a complex technique involving both histological and immunological detection methods. It requires a highly trained histotechnologist. Tissue processing and handling prior to immunostaining, for example variations in fixation and embedding or the inherent nature of the tissue can cause inconsistent results (Nadji and Morales, 1983). Inadequate counterstaining and mounting can influence the interpretation of the results. Monosan® guarantees that the product will meet all requirements described from its shipping date until its expiry date, as long as the product is correctly stored and utilized. No additional guarantees can be given. Under no circumstances shall Monosan® be liable for any damages arising out of the use of the reagent provided.

## Performance characteristics










Monosan® has conducted studies to evaluate the performance of the reagent. The product has been found to be suitable for the intended use.

## Bibliography

Varma M, Linden MD, Amin MB Mod Pathol 12:472-478, 1999  
Nadji M and Morales AR Ann N.Y. Acad Sci 420:134-9, 1983

FOR RESEARCH USE ONLY, NOT FOR DRUG, DIAGNOSTIC OR OTHER USE.

Explanation of the symbols on the product label:

 REF	Bestellnummer Catalog Number Reference du catalogue	 LOT	Chargenbezeichnung Batch Code Code du lot	 X <sub>n</sub>	Reizend Irritant Irritant
 X <sub>n</sub>	Gesundheitsschädlich Harmful Nocif		Giftig Toxic Toxique	 Hersteller / Manufacturer / Fabricant  Monosan® Frontstraat 2c 5405 PB Uden The Netherlands Tel: (+31) 413 251115 Fax: (+31) 413 266605 info@monosan.com www.monosan.com	
	Verwendbar bis Use By Utiliser jusque				
	Gebrauchsanweisung beachten Consult Instructions for use Consulter les instructions d'utilisation		Lagerungstemperatur Temperature Limitation Limites de température		