



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-BAP1, clone C-4 (Monoclonal)

Clone no. C-4

MONOSAN

Product name	Mouse anti-BAP1, clone C-4 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:50-1:500), WB (1:100-1:500)
Species reactivity	Human
Conjugate	-
Immunogen	Amino acids 430-729 of BAP1 of human origin
Isotype	IgG1
Clonality	Monoclonal
Clone number	C-4
Size	1 ml
Concentration	n/a
Format	Concentrate
Storage buffer	PBS with < 0.1% sodium azide and 0.1% gelatin
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-BAP1, clone C-4 (Monoclonal)

Clone no. C-4

MONOSAN

Additional info

Mutations within the BRCA1 gene, localized to chromosome 17q, are believed to account for approximately 45% of families with increased incidence of both early-onset breast cancer and ovarian cancer. The BRCA1 gene is expressed in numerous tissues, including breast and ovary, and encodes a predicted protein of 1,863 amino acids. This protein contains a RING domain near the N-terminus and appears to encode a tumor suppressor. BARD1 (BRCA1-associated RING domain protein 1) and BAP1 (BRCA1-associated protein 1) have both been shown to bind to the N-terminus of BRCA1 and are potential mediators of tumor suppression. BARD1 contains an N-terminal RING domain and three tandem ankyrin repeats. The C-terminus of BARD1 contains a region with sequence homology to BRCA1, termed the BRCT domain. BAP1 is a ubiquitin hydrolase and has been shown to enhance BRCA1-mediated cell growth suppression. Pre-treatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, or in 50 mM Tris buffer pH9.5, for 20 minutes is required for IHC staining on formalin-fixed, paraffin embedded tissue sections. Control tissue Pancreas, breast carcinoma, ovarian carcinoma. Staining Nuclear and cytoplasmic.

References

1. Hall, J.M., et al. 1990, Science 250: 1684-1689.
2. Yoshikawa, Y., et al. 2012, Cancer Sci. 103: 868-874.
3. Gammon, B., et al. 2013, J. Cutan. Pathol. 40: 538-542.
4. Kerl, K., et al. 2013, Am. J. Dermatopathol. 35: 151-158.
5. Popova T., et al. 2013, Am. J. Hum. Genet. 92: 974-980.

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES