



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-YAP, clone 63.7 (Monoclonal)

Clone no. 63.7

MONOSAN

Product name	Mouse anti-YAP, clone 63.7 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:500-1:1000)
Species reactivity	Human
Conjugate	-
Immunogen	Recombinant YAP of human origin
Isotype	IgG2a
Clonality	Monoclonal
Clone number	63.7
Size	1 ml
Concentration	n/a
Format	Concentrate
Storage buffer	PBS with less than 0.1% sodium azide and 0.1% gelatin
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-YAP, clone 63.7 (Monoclonal)

Clone no. 63.7

MONOSAN

Additional info

The Yes-associated protein, otherwise known as YAP, is a 14-3-3 binding molecule that was originally recognized by virtue of its ability to bind to the SH3 domain of Yes. The binding of YAP to 14-3-3 requires the phosphorylation of a homologous serine residue (Ser 112) in the YAP 14-3-3 binding motif. The highly conserved and ubiquitously expressed 14-3-3 proteins regulate differentiation, cell cycle progression and apoptosis by binding intracellular phosphoproteins involved in signal transduction. YAP may link events at the plasma membrane and cytoskeleton to inhibition of transcription in the nucleus in a manner regulated by 14-3-3 proteins. YAP shares homology with the WW domain of TAZ, transcriptional co-activator with PDZ binding motif, which functions as a transcriptional co-activator by binding to the PPXY motif present in transcription factors. YAP is expressed at high levels in the lung, placenta, prostate, ovary and testis. Pretreatment: Heat induced epitope retrieval in 10 mM citrate buffer, pH6.0, for 20 minutes is required for IHC staining on formalin-fixed, paraffin embedded tissue sections. Note: Dilution of the antibody in 10% normal goat serum followed by a goat anti-mouse secondary antibody-based detection is recommended. Control tissue Lung, placenta, prostate, ovary, testis. Staining Nuclear.

References

1. Ren, Y.R., et al. 2011. S, J. Biol. Chem. 286: 11960-11969.
2. Ellison, D.W., et al. 2011. Acta Neuropathol. 121: 381-396.
3. Cordenonsi, M., et al. 2011. Cell 147: 759-772.
4. Ren, Y.R., et al. 2012. J. Proteome Res. 11: 5301-5310.
5. Vigneron, A.M. and Vousden, K.H. 2012. EMBO J. 31: 471-480.

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES