



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Mouse anti-Human MMP-9 Activated, clone 4A3 (Monoclonal)

Clone no. 4A3

MONOSAN

---

|                           |   |
|---------------------------|---|
| Product name              | Mouse anti-Human MMP-9 Activated, clone 4A3 (Monoclonal)  |
| Host                      | Mouse   |
| Applications              | IHC-P, WB   |
| Species reactivity        | human   |
| Conjugate                 | -   |
| Immunogen                 | Ovalbumin conjugated synthetic peptide corresponding to a region within the N-terminus of human MMP-9 |
| Isotype                   | IgG1  |
| Clonality                 | Monoclonal  |
| Clone number              | 4A3   |
| Size                      | 0.2 mg  |
| Concentration             | IgG 0.5 mg/ml   |
| Format                    | Purified  |
| Storage buffer            | PBS with <0.1% sodium azide   |
| Storage until expiry date | aliquots -20°C. Thawed 2-8°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti-Human MMP-9 Activated, clone 4A3 (Monoclonal)

Clone no. 4A3

MONOSAN

**Additional info**

Mouse anti Human MMP-9 Activated antibody, clone 4A3 recognizes the active form of human matrix metalloproteinase 9 (MMP-9). The MMP's are zinc-dependent endopeptidases responsible for degrading the extracellular matrix. They are also involved in cell proliferation, migration, differentiation and apoptosis. Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active. Their expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis. MMP-9 is a gelatinase, cleaving type IV collagen and gelatin. The gelatinases have an additional gelatin-binding domain inserted in the catalytic domain. MMP-9, alongside other MMPs, plays a role in normal tissue remodeling such as embryonic development, ovulation, mammary gland involution and wound healing. It is important in the early stages of tumor invasion as it degrades the type IV collagen in the basement membrane

**References**

1. Duncan ME et al. Eur J Biochem 1998; 258: 37-43
2. Selemetjev S et al. Am J Clin Pathol 2016; pii: aqw184 (Epub ahead of print)
3. -
4. -
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**