



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Human Von Willebrand Factor, clone RFF-VIII R/1 (Monoclonal)

Clone no. RFF-VIII R/1

MONOSAN

Product name	Mouse anti-Human Von Willebrand Factor, clone RFF-VIII R/1 (Monoclonal)
Host	Mouse
Applications	IHC-fr, ELISA
Species reactivity	human
Conjugate	-
Immunogen	Human Factor VIII complex partially purified from Factor VIII concentrate
Isotype	IgG1
Clonality	Monoclonal
Clone number	RFF-VIII R/1
Size	0.2 mg
Concentration	IgG 1 mg/ml
Format	Purified
Storage buffer	PBS with <0.1% sodium azide
Storage until expiry date	aliquots -20°C. Thawed 2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Human Von Willebrand Factor, clone RFF-VIII R/1 (Monoclonal)

Clone no. RFF-VIII R/1

MONOSAN

Additional info

Mouse anti Human von Willebrand factor antibody, clone RFF-VIII R/1 recognizes human von Willebrand factor (vWF), also known as Factor VIII related antigen, a blood glycoprotein involved in blood coagulation. It stabilises circulating Factor VIII by binding to it and protecting it from cleavage and delivers it to sites of vascular injury. vWF also promotes the adhesion of platelets to sites of vascular damage by forming a molecular bridge between collagen on exposed endothelial cells and the GPIb binding sites of platelets circulating in the blood. vWF circulates in the blood as large multimers, with each monomer (250kDa) containing a number of specific domains. Mouse anti Human von Willebrand factor, clone RFF-VIII R/1 has a high affinity for an epitope within the platelet GPIb-binding site that is responsible for biological activity. As such the antibody is a potent inhibitor of vWF activity. It can completely neutralise ristocetin-induced platelet aggregation and ristocetin-induced binding of vWF to platelets. It also inhibits platelet adhesion to glass beads. The epitope recognized is present only on the intact multimeric form of vWF and is abolished by mild denaturation with SDS. Mouse anti Human von Willebrand factor, clone RFF-VIII R/1 does not recognize human Factor VIII.

References

1. Goodall AH et al. Br J Haematol 1985; 59: 565-77
2. Kraus E et al. Thromb Res 2014; 2134: 1285-91
3. Chen YJ et al. Stroke 2015; 46: 237-44
4. Tejada de Rink MM et al. Ther Hypothermia Tem Manag 2020; 10: 91-101
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES