



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal)

Clone no. SQab1880

MONOSAN

Product name	Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal)
Host	Rabbit
Applications	FC, ICC/IF, IP, WB
Species reactivity	Other
Conjugate	-
Immunogen	KLH-conjugated DYKDDDDK.
Isotype	-
Clonality	Monoclonal
Clone number	SQab1880
Size	100 ul
Concentration	n/a
Format	Purification with Protein A.
Storage buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.
Storage until expiry date	-20°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-DDDDK tag antibody, clone SQab1880 (monoclonal)


Clone no. SQab1880

MONOSAN


Additional info

Application note: * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background:

Images



Immunofluorescence: 293 cells transfected with C-terminal DYKDDDDK tagged gene. Cells were fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% goat serum for 30 min at RT. Cells were stained with DAPI (left)



Western blot: Testing of anti-DDDDK tag antibody [SQab1880]. 3 µg of 293 cell lysate without any transfection (at 1:2000, left) or transfected with N-terminal DYKDDDDK tagged gene (at 1:10000, right).

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES