



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Rabbit anti-V5 tag antibody, clone SQab18114 (monoclonal)

Clone no. SQab18114

MONOSAN

| | |
|---------------------------|---|
| Product name | Rabbit anti-V5 tag antibody, clone SQab18114 (monoclonal) |
| Host | Rabbit |
| Applications | FC, ICC/IF, IP, WB |
| Species reactivity | Human, Mouse, Rat |
| Conjugate | - |
| Immunogen | GKPIPPLLGLDST (V5 epitope) conjugated to KLH. |
| Isotype | - |
| Clonality | Monoclonal |
| Clone number | SQab18114 |
| Size | 100 ul |
| Concentration | n/a |
| Format | Purification with Protein A. |
| Storage buffer | PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA. |
| Storage until expiry date | -20°C |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Product datasheet

MON240177

MONOSAN[®]

Rabbit anti-V5 tag antibody, clone SQab18114 (monoclonal)


Clone no. SQab18114

MONOSAN


Additional info

Application note: * The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background:

Images



Immunofluorescence: 293 cells transfected with N-terminal V5 tagged gene. Cells were fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% goat serum for half an hour at room temperature. Samples were



Western blot: 2 µg of 1) 293 cells transfected with N-terminal V5 tagged gene and stained with anti-V5 tag antibody [SQab18114] at 1:10000 dilution. 2) 293 cells transfected with C-terminal V5 tagged gene and stained with anti-V5 tag antibody [SQab18114] at 1:2000 dilution. 3) Mock 293

References

- 1. -
- 2. -
- 3. -
- 4. -
- 5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES