



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rabbit anti-HA tag antibody, clone SQab18117 (monoclonal)

Clone no. SQab18117

MONOSAN

---

Product name	Rabbit anti-HA tag antibody, clone SQab18117 (monoclonal)
Host	Rabbit
Applications	FC, ICC/IF, IP, WB
Species reactivity	Human, Mouse, Rat, African green monkey, Bovine, Chicken, Dog
Conjugate	-
Immunogen	YPYDVPDYA (influenza hemagglutinin-HA-epitope) conjugated to KLH.
Isotype	-
Clonality	Monoclonal
Clone number	SQab18117
Size	100 ul
Concentration	n/a
Format	Purification with Protein A.
Storage buffer	PBS, 0.01% Sodium azide, 40% Glycerol and 0.05% BSA.
Storage until expiry date	-20°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Product datasheet

MON240179

**MONOSAN**<sup>®</sup>

Rabbit anti-HA tag antibody, clone SQab18117 (monoclonal)

Clone no. SQab18117


MONOSAN

---


#### Additional info

Application note: \* The dilutions indicate recommended starting dilutions and the optimal dilutions or concentrations should be determined by the scientist. Storage instruction: For continuous use, store undiluted antibody at 2-8°C for up to a week. For long-term storage, aliquot and store at -20°C. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening. The antibody solution should be gently mixed before use. Background:

Images



Immunofluorescence: 293 cells transfected with N-terminal HA tagged gene. Cells were fixed with 4% paraformaldehyde for 30 min at RT, permeabilized with 0.1% Triton X-100 for 10 min at RT then blocked with 10% Goat serum for half an hour at room temperature. Samples were



Immunoprecipitation: 0.2 mg of 293 whole cell lysate (transfected with N-terminal HA tagged gene) was immunoprecipitated (1:10 dilution) and stained with anti-HA tag antibody [SQab18117].

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES