



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Mouse anti-Epithelial Keratin Type I, clone AE-1 (Monoclonal)

Clone no. AE-1

MONOSAN

Product name Mouse anti-Epithelial Keratin Type I, clone AE-1 (Monoclonal)

Host Mouse

Applications IHC-P (1:150-1:200), IHC-Fr (1:150-1:200), WB (1:100)

Species reactivity human, mouse, rabbit, rat, bovine, chicken

Conjugate -

Immunogen Human epidermal keratin

Isotype IgG1

Clonality Monoclonal

Clone number AE-1

Size 250 ug

Concentration 1 mg/ml

Format -

Storage buffer PBS with 0.1% sodium azide

Storage until expiry date 2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Epithelial Keratin Type I, clone AE-1 (Monoclonal)

Clone no. AE-1

MONOSAN

Additional info

Cytokeratin pan is part of a subfamily of intermediate filament proteins that are characterized by remarkable biochemical diversity, and represented in human epithelial tissues by at least 20 different polypeptides. Cytokeratins range in molecular weight between 40 kDa- 68 kDa, and an isoelectric pH between 4.9-7.8. The individual human cytokeratins are numbered 1 to 20. The various epithelia in the human body usually express cytokeratins which are not only characteristic of the type of epithelium, but also related to the degree of maturation or differentiation within an epithelium. Cytokeratin subtype expression patterns are used to an increasing extent in the distinction of different types of epithelial malignancies. The cytokeratin antibodies are not only of assistance in the differential diagnosis of tumors using immunohistochemistry on tissue sections, but are also a useful tool in cytopathology and flow cytometric assays. The composition of cytokeratin pairs vary with the epithelial cell type, stage of differentiation, cellular growth environment, and disease state. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis.

References

1. Woodcock-Mitchell, J., et al., J. Cell Biol. 95, 580-588 (1982).
2. Tseng, S.C.G., et al., Cell 30, 361-372 (1982).
3. Eichner, R., et al., J. Cell Biol. 98, 1388-1396 (1984)
4. Sun, T.-T., et al., The Cancer Cell 1,169-176 (1984)
5. Battifora, H., et al., Cancer 54, 843-848 (1984)

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES