



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Cardiotin, clone SR-2, Purified (Monoclonal)

Clone no. SR-2

MONOSAN

Product name	Mouse anti-Cardiotin, clone SR-2, Purified (Monoclonal)
Host	Mouse
Applications	IHC-fr (1:25-1:100), IHC-P (1:25-1:100), WB (1:50-1:500)
Species reactivity	human, swine
Conjugate	-
Immunogen	100 kDa cardiotin subunit
Isotype	IgM
Clonality	Monoclonal
Clone number	SR-2
Size	0.1 mg
Concentration	n/a
Format	-
Storage buffer	PBS with 0.09% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Cardiotin, clone SR-2, Purified (Monoclonal)

Clone no. SR-2

MONOSAN

Additional info

Cardiotin is a high molecular weight protein complex (300 kDa) located in the mitochondria of cardiomyocytes and skeletal muscle. The cardiotin structure exists of subunits of 60 kDa and 100 kDa, probably in a tetrameric configuration. Both subunits contain the same amino-terminal 14 amino-acid sequence, showing high homology to Human skeletal muscle α -actinin. During cardiac contractile dysfunction and myocard cell differentiation, the cardiotin distribution is affected. Compared to other structural proteins, cardiotin is one of the first to respond to insults (ischemia, fibrillation) that influence the functional status of cardiomyocytes. SR-2 reacts with cardiomyocytes, skeletal muscle, stromal and epithelial cells as well in vivo as in vitro.

References

1. Schaart et al. FEBS Lett;403;168-172
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES