



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Mouse anti-Lewis B, clone 2-25LE (Monoclonal)

Clone no. 2-25LE

MONOSAN

---

|                           |   |
|---------------------------|---|
| Product name              | Mouse anti-Lewis B, clone 2-25LE (Monoclonal) |
| Host                      | Mouse   |
| Applications              | ELISA, FC, IHC-fr, IHC-P, IF, WB              |
| Species reactivity        | human, guinea pig                             |
| Conjugate                 | -   |
| Immunogen                 | ALea-b+ ovarian cyst mucin                    |
| Isotype                   | IgG1  |
| Clonality                 | Monoclonal                                    |
| Clone number              | 2-25LE  |
| Size                      | 100 ug  |
| Concentration             | 100 ug/ml                                     |
| Format                    | -   |
| Storage buffer            | PBS with 0.02% sodium azide                   |
| Storage until expiry date | 2-8°C   |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Lewis B, clone 2-25LE (Monoclonal)

Clone no. 2-25LE

MONOSAN

---

**Additional info**

2-25LE reacts with Lewis b antigen which is highly expressed in stomach, colon, small intestine, lung and kidney and to a lesser extent in salivary gland, bladder, uterus and liver

**References**

1. Bara et al. Cancer Res. 46: 3983-3898 (1986)
2. Bara et al. Biochem. J. 254: 185-193 (1988)
3. Creuzot-Garcher et al. Invest Oph(thalmol Vis Sci. 40(8):1631-6 (1999)
4. Good et al. Vox Sanguinis 62(3): 180-189 (1992)
5. Torrado et al. Gastroenterology, 102 (2): 424-430 (1992)

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES