



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rat anti-Mouse F4/80, clone BM8 (Monoclonal)

Clone no. BM8

MONOSAN

Product name	Rat anti-Mouse F4/80, clone BM8 (Monoclonal)
Host	Rat
Applications	IHC-fr,FC,IHC-P,WB
Species reactivity	mouse
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG2a
Clonality	Monoclonal
Clone number	BM8
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rat anti-Mouse F4/80, clone BM8 (Monoclonal)

Clone no. BM8

MONOSAN

Additional info

The monoclonal antibody BM8 recognises a 125 kD extracellular macrophage membrane molecule, highly restricted to mature macrophage subpopulations residing in tissue. This murine F4/80 glycoprotein contains seven-transmembrane (TM7) regions, which anchor the protein in the cell membrane, and thereby shows similarity in this region to G-protein-coupled receptors. The F4/80 molecule shares overall structural homology to other members of the epidermal growth factor (EGF)-TM7 family. The antigen is detected on tissue fixed macrophages in all organs tested so far (spleen, lymph nodes, thymus, liver, skin). It is also present on Langerhans cells in the skin and Kupffer cells in the liver. It is absent on granulocytes, lymphocytes and trombocytes. The expression of F4/80 increases upon maturation of macrophage precursors in bone marrow and blood as well as in ontogeny.

The monoclonal antibody BM8 is the only macrophage marker that is able to distinguish non-destructive from destructive inflammation processes in the pancreas. Furthermore it is a unique histological marker of the progression from peri-insulinitis to beta-cell destruction and diabetes in a mouse diabetes model.

References

1. Malorny; U et al. Cell Tissue Res 1986; 243: 421
2. Leenen, P et al J Immunol Methods 1994, 174: 5
3. Mackler; A et al. Biol Reprod 1999; 61: 879
4. Schaller E et al. Mol Cell Biol 2002; 22: 8035
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES