



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD10, clone 56C6 (Monoclonal)

Clone no. 56C6

MONOSAN

---

Product name	Mouse anti-CD10, clone 56C6 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:10-1:50)
Species reactivity	human
Conjugate	-
Immunogen	Recombinant protein encoding external domain of human CD10
Isotype	IgG1
Clonality	Monoclonal
Clone number	56C6
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD10, clone 56C6 (Monoclonal)

Clone no. 56C6

MONOSAN

**Additional info**

Common acute lymphoblastic leukemia antigen (CALLA / CD10) is a useful marker for the characterization of childhood leukemia and B cell lymphomas. This antibody reacts with antigen of lymphoblastic, Burkitt's, and follicular lymphomas; and chronic myelocytic leukemia. Also, Anti-CD10 detects the antigen of glomerular epithelial cells and the brush border of the proximal tubules; this characteristic may be helpful in interpreting renal ontogenesis in conjunction with other markers. Other non-lymphoid cells that are reactive with CD10 are breast myoepithelial cells, bile canaliculi, neutrophils and small population of bone marrow cells, fetal small intestine epithelium, and normal fibroblasts.

**References**

1. Maguer-Satta V, et al.. Stem Cells. 2011; 29:389-96
2. Laurent C, et al. 2010; 95: 356-8
3. Onak NK, et al.. Turk J Med Sci. 2010; 40:177-84
4. Chu P, et al. Am J Clin Pathol. 2000; 113: 374-82
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**