



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD7, clone MRQ-56 (Monoclonal)

Clone no. MRQ-56

MONOSAN

Product name	Mouse anti-CD7, clone MRQ-56 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:100-1:500)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1-k
Clonality	Monoclonal
Clone number	MRQ-56
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD7, clone MRQ-56 (Monoclonal)

Clone no. MRQ-56

MONOSAN

Additional info

CD7 antigen is a 40-kDa cell surface glycoprotein that is a member of the immunoglobulin gene superfamily. While its precise function is not known, it is suggested that CD7 plays a role in T-cell interactions as it is one of the earliest T-cell lineage associated antigens expressed during T-cell ontogeny. CD7 is expressed in thymocytes, mature peripheral T-cells, natural killer cells, and lymphoid and myeloid progenitors. CD7 is the most consistently expressed T cell antigen in lymphoblastic lymphomas and leukemias, and is therefore a useful marker in the identification of such neoplastic proliferations. In mature post-thymic T cell neoplasms, it is the most common pan-T antigen to be aberrantly absent and its absence in a T cell population is a useful pointer to a neoplastic conversion.

References

1. Hodak E, et al. J Am Acad Dermatol. 2006 Aug;55(2):276-84
2. Stillwell R, et al. Immunol Res. 2001; 24:31-52
3. Schanberg LE, et al. Proc Natl Acad Sci USA. 1991; 88:603-7
4. Chabannon C, et al. J Immunol. 1992; 149:2110-3
5. Rabinowich H, et al. J Immunol. 2003; 152: 517-26

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES