



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Mouse anti-Factor XIIIa, clone AC-1A1 (Monoclonal)

Clone no. AC-1A1

MONOSAN

---

|                           |   |
|---------------------------|---|
| Product name              | Mouse anti-Factor XIIIa, clone AC-1A1 (Monoclonal)                |
| Host                      | Mouse   |
| Applications              | IHC-P (1:100-1:500)   |
| Species reactivity        | human   |
| Conjugate                 | -   |
| Immunogen                 | Unknown or proprietary to MONOSAN and/or its suppliers            |
| Isotype                   | IgG1  |
| Clonality                 | Monoclonal  |
| Clone number              | AC-1A1  |
| Size                      | 1 ml  |
| Concentration             | n/a   |
| Format                    | -   |
| Storage buffer            | Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide |
| Storage until expiry date | 2-8°C   |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Factor XIIIa, clone AC-1A1 (Monoclonal)

Clone no. AC-1A1

MONOSAN

**Additional info**

Factor XIIIa has been identified in platelets, megakaryocytes, and fibroblast-like mesenchymal or histiocytic cells in the placenta, uterus, and prostate, monocytes and macrophages and dermal dendritic cells. Anti-factor XIIIa has been found to be useful in differentiating between dermatofibroma (almost all cases +), dermatofibrosarcoma protuberans (-/+) and desmoplastic malignant melanoma. Anti-factor XIIIa positivity is also seen in capillary hemangioblastoma, hemangioendothelioma, hemangiopericytoma, xanthogranuloma, xanthoma, hepatocellular carcinoma, glomus tumor, and meningioma.

**References**

1. Nemes Z, Hum Pathol 1992 Jul; 23(7):805-10
2. Horenstein MG et al. Am J Surg Pathol. 2000 Jul;24(7):996-1003
3. Kraus MD et al. Am J Dermatopathol. 2001 Apr;23(2):104-11
4. Dehner LP. Am J Surg Pathol. 2003 May;27(5):579-93
5. Deguchi M et al. Arch Dermatol Res. 2002 Oct;294(7):297-302

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**