



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Human MUC1, clone MRQ-17 (Monoclonal)

Clone no. MRQ-17

MONOSAN

Product name	Mouse anti-Human MUC1, clone MRQ-17 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:50-1:200)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	MRQ-17
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Human MUC1, clone MRQ-17 (Monoclonal)

Clone no. MRQ-17

MONOSAN

Additional info

Mucins are high molecular weight glycoproteins which constitute the major component of the mucus layer that protects the gastric epithelium from chemical and mechanical injury. In humans, at least 14 mucin genes have been identified that code for the mucin proteins. Mucin genes are expressed in a regulated cell- and tissue-specific manner. The stomach provides a good example of such differential expression of mucin genes. MUC1 is detected in mucous cells of the surface epithelium and neck region of the gastric antrum, as well as in pyloric glands and oxyntic glands of the body region. The heterogeneous pattern of mucin expression, including the expression of the intestinal mucin MUC2, may provide new insights into the differentiation pathways of gastric carcinoma.

References

1. Chaves P, et al. Dis Esophagus. 2005; 18:383-7
2. Leteurtre E, et al. World J Gastroenterol. 2006; 12:3324-31.
3. Mino-Kenudson M, et al. Arch Pathol Lab Med. 2007; 131:86-90
4. Mizoshita T, et al. Histol Histopathol. 2007; 22:251-60
5. O'Connell FP, et al. Arch Pathol Lab Med. 2005; 129:338-47

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES