



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rabbit anti-Human T-bet, clone MRQ-46 (Monoclonal)

Clone no. MRQ-46

MONOSAN

Product name	Rabbit anti-Human T-bet, clone MRQ-46 (Monoclonal)
Host	Rabbit
Applications	IHC-P (1:100-1:500)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	MRQ-46
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-Human T-bet, clone MRQ-46 (Monoclonal)

Clone no. MRQ-46

MONOSAN

Additional info

T-bet, a T-box transcription factor, is expressed in CD4+ T-lymphocytes committed to T-helper (Th)1 Tcell development from naïve T-helper precursor cells (Thp) and redirects Th2 T-cells to Th1 development. Anti-T-bet is a marker of mature T-cells and is expressed at very low levels in Thp cells and is absent in precursor T-lymphoblastic leukemia/lymphoma cells. Scattered small lymphocytes in the interfollicular T-cell zone of reactive lymphoid tissue, including tonsil, lymph node, and spleen exhibited nuclear staining for anti-T-bet, with no anti-T-bet staining observed in germinal centers or mantle or marginal zones. T-bet is expressed in a significant subset of B-cell lymphoproliferative disorders, particularly at an early stage of B-cell development (precursor B-cell lymphoblastic leukemia/lymphoblastic lymphoma), and B-cell neoplasms derived from mature B-cells, including CLL/SLL, marginal zone lymphoma, and hairy cell leukemia.

References

1. Szabo SJ, et al. Cell. 2000; 100:665-69
2. Jöhrens K, et al. Am J Surg Pathol. 2007; 31:1181-5
3. Atayar C, et al. Am J Pathol. 2005; 166:127-34
4. Dorfman DM, et al. Am J Clin Pathol. 2004; 122:292-7
5. Harashima A, et al.. Leuk Res. 2005; 29:841-8

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES