



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-MadCAM-1, clone 314G8 (Monoclonal)

Clone no. 314G8

MONOSAN

Product name	Mouse anti-MadCAM-1, clone 314G8 (Monoclonal)
Host	Mouse
Applications	IHC-fr,FUNC,ELISA,IHC-P,WB
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	314G8
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-MadCAM-1, clone 314G8 (Monoclonal)

Clone no. 314G8

MONOSAN

Additional info

The monoclonal antibody 314G8 reacts with human mucosal addressin cell adhesion molecules-1 (MAdCAM-1), a key player in mediating the infiltration of leukocytes into chronically inflamed tissue. MAdCAM-1 is a cell-surface Ig superfamily member composed of two extracellular Ig domains, followed by a mucin-like domain, a transmembrane domain and a short cytoplasmic domain. It interacts via its N-terminal Ig domain with the lymphocyte homing receptor alpha4beta7, which plays a critical role in forming the gut-associated lymphoid system. MAdCAM-1 promotes the adhesion of T- and B cells, monocytes/macrophages, and potentially eosinophils, basophils, and differentiated mast cells to the vascular endothelium. Mucosal addressin cell adhesion molecule-1 RNA transcripts are predominantly expressed in the small intestine, mesenteric lymph nodes, colon and spleen; and are very weakly expressed in human pancreas and brain. The monoclonal antibody 314G8 recognizes a site in the N-terminal Ig domain of MAdCAM-1. The monoclonal antibody 314G8 detects MAdCAM-1 on venules in the spleen and small intestine. MAdCAM-1 is strongly expressed in the synovium of osteoarthritis patients, predominantly on the endothelial lining of blood vessels, but also within the vessel lumen. The monoclonal antibody 314G8 may be useful in diagnosis of inflammation in humans by monitoring the presence and levels of MAdCAM-1.

References

1. Leung; E et al. Immunol Cell Biol 2004; 82: 400
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES