



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-TLR2, clone TL2.3 (Monoclonal)

Clone no. TL2.3

MONOSAN

Product name	Mouse anti-TLR2, clone TL2.3 (Monoclonal)
Host	Mouse
Applications	IHC-fr,FC,FUNC,ELISA,IF,WB
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG2a
Clonality	Monoclonal
Clone number	TL2.3
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-TLR2, clone TL2.3 (Monoclonal)

Clone no. TL2.3

MONOSAN

Additional info

Toll-like receptors (TLR) are highly conserved throughout evolution and have been implicated in the innate defense to many pathogens. In *Drosophila*, toll is required for the anti-fungal response, while the related 18-wheeler is involved in antibacterial defenses. In mammals, TLR identified as type I transmembrane signaling receptors with pattern recognition capabilities, have been implicated in the innate host defense to pathogens. TLR2 has been identified as a receptor that is central to the innate immune response to lipoproteins of Gram-negative bacteria, several whole Gram-positive bacteria, as well as a receptor for peptidoglycan and lipoteichoic acid and other bacterial cell membrane products. A functional interaction between TLR2 and TLR6 in the cellular response to various bacterial products has been discovered. The currently accepted paradigm regards TLR2 as an essential receptor for many eubacterial cell wall components, including lipoproteins and peptidoglycan. Bacterial species as diverse as mycobacteria, spirochetes, mycoplasma, *Staphylococcus aureus*, and *Streptococcus pneumoniae* have all been shown to mediate cellular activation via TLR2 (CD282). The monoclonal antibody TL2.3 is specific for human TLR2 (CD282). TL2.3 is useful for studies on the role of TLR2 as a pattern recognition receptor in microbial products induced cytokine production by TLR2 bearing cells such as human peripheral blood mononuclear cells. The monoclonal antibody TL23 is cross reactive with canine TLR2.

References

1. Flo; T et al. *J Leukoc Biol* 2001; 69: 474
2. Schjetne, K et al *J Immunol* 2003, 171: 32
3. Siedlar; M et al. *J Immunol* 2004; 173: 2736
4. Tunheim G et al. *Vaccine* 2007; 25: 4723
5. Burgener I et al. *Vet Immunol Immunopathol* 2008; 124: 184

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES