



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product datasheet MON4043

MONOSAN[®]

Mouse anti-CD130, clone B-T2 (Monoclonal)

Clone no. B-T2

MONOSAN

Product name Mouse anti-CD130, clone B-T2 (Monoclonal)

Host Mouse

Applications ELISA,FC

Species reactivity human

Conjugate Purified

Immunogen Natural soluble gp130 receptor.

Isotype IgG1

Clonality Monoclonal

Clone number B-T2

Size 0.1 mg

Concentration 1 mg/ml

Format -

Storage buffer PBS

Storage until expiry date 2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD130, clone B-T2 (Monoclonal)

Clone no. B-T2

MONOSAN

Additional info

Mouse anti Human CD130 antibody, clone B-T2 recognizes soluble and membrane bound gp130 receptor, also known as CD130, Interleukin-6 receptor subunit beta, Oncostatin-M receptor subunit alpha or Interleukin-6 signal transducer. CD130 is a 918 amino acid, 130 kDa single pass type-1 transmembrane glycoprotein containing a single Ig-like C2 type and multiple fibronectin type-III domains. CD130 can be cleaved to form a monomeric soluble 100 kDa form of the receptor detectable in plasma and other biologic fluids where it acts as an IL-6 antagonist (Miller-Newen et al. 1998). Mouse anti Human CD130 antibody, clone B-T2 binds to an epitope dependent at least partially on the presence of the Ig-like domain (Hammacher et al. 1998). Mouse anti Human CD130 antibody, clone B-T2 has been reported to block receptor activity and inhibit IL-6 induced proliferation of XG1 cells and IL-27 mediated proliferation of naive CD4+ T cells (de Groot et al. 2012, Pflanz et al. 2004).

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES