



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD91, clone A2Mr alpha-2 (Monoclonal)

Clone no. A2Mr

MONOSAN

Product name	Mouse anti-CD91, clone A2Mr alpha-2 (Monoclonal)
Host	Mouse
Applications	WB,IP,FC,IHC-fr,ELISA
Species reactivity	human
Conjugate	Purified
Immunogen	Purified alpha2 macroglobulin receptor
Isotype	IgG1
Clonality	Monoclonal
Clone number	A2Mr alpha-2
Size	0.1 mg
Concentration	1.0mg/ml
Format	-
Storage buffer	PBS with 0.09% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD91, clone A2Mr alpha-2 (Monoclonal)

Clone no. A2Mr alpha-2

MONOSAN

Additional info

Mouse anti Human CD91 antibody, clone A2Mr alpha-2 recognizes human CD91, also known as Prolow-density lipoprotein receptor-related protein 1, Alpha-2-macroglobulin receptor or apolipoprotein E receptor. CD91 is a 4525 amino acid protein post translationally cleaved into 3 subunits, a 85 kDa type I transmembrane carboxyl chain (LRP85) non-covalently bound to a 515 kDa extracellular N-terminal subunit (LRP515) containing multiple EGF-like and LDL-receptor Class A and Class B domains. Additionally, there is an intracellular domain (LRPICD) which can be cleaved from the transmembrane domain by gamma secretase (May et al. 2004). Clone A2Mr alpha-2 detects an epitope within the LRP515 chain. CD91 is a multifunctional protein involved in processes including the phagocytosis and endocytosis of apoptotic cells (Nilsson et al. 2012), clearance of activated serum alpha-2-macroglobulin (Kristensen et al. 1990), modulation of the inflammatory response (Staudt et al. 2013) and acts as a receptor for Pseudomonas aeruginosa exotoxin A (Kounnas et al. 1992). Mouse anti Human CD91, clone A2Mr alpha-2 has been used extensively for the detection of CD91 by flow cytometry and immunohistochemistry

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES