



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-CD229, clone HLy 9.1.25 (Monoclonal)

Clone no. HLy9.1.25

MONOSAN

Product name	Mouse anti-CD229, clone HLy 9.1.25 (Monoclonal)
Host	Mouse
Applications	IP,FC
Species reactivity	human
Conjugate	Purified
Immunogen	Mouse pre-B cell line 300.19 transfected with CD229 cDNA.
Isotype	IgG1
Clonality	Monoclonal
Clone number	HLy9.1.25
Size	0.2 mg
Concentration	1.0 mg/ml
Format	-
Storage buffer	PBS with 0.09% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD229, clone HLy 9.1.25 (Monoclonal)

Clone no. HLy9.1.25

MONOSAN

Additional info

Mouse anti Human CD229 antibody, clone HLy9.1.25 recognizes the human cell surface antigen CD229 also known as T-lymphocyte surface antigen Ly-9 or SLAM family member 3. CD229 is a 608 amino acid single pass type I transmembrane glycoprotein of ~126;120 kDa as evaluated by immunoprecipitation of cells transfected with the full length human CD229 cDNA. However immunoprecipitation of CD229 from Daudi cell lysates with clone HLy9.1.25 yields bands of 120 kDa corresponding to the full length CD229 and a ~126;100 kDa band attributed to an alternatively spliced isoform lacking the fourth Ig-like domain (de la Fuente et al. 2001). Human CD229 is expressed on thymocytes, T-cells and B-cells (Del Valle et al. 2003). CD229 has also been described as a tumor associated antigen in chronic lymphocytic leukemia (Bund et al. 2006) and has been implicated in the development of spontaneous autoantibody production to nuclear antigens in mice and is potentially a target for the treatment of autoimmunity (de Salort et al. 2013).

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES