



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Tumor Necrosis factor Alpha, clone 52B83 (Monoclonal)

Clone no. 52B83

MONOSAN

Product name	Mouse anti-Tumor Necrosis factor Alpha, clone 52B83 (Monoclonal)
Host	Mouse
Applications	IHC-fr,FC,ELISA,WB
Species reactivity	human, guinea pig, mouse, rhesus monkey
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	52B83
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Tumor Necrosis factor Alpha, clone 52B83 (Monoclonal)

Clone no. 52B83

MONOSAN

Additional info

The monoclonal antibody 52B83 reacts with tumor necrosis factor alpha (TNF-alpha). TNF-alpha is a homotrimeric 17 kDa protein, that interacts with either one of the two types of TNF-receptors, termed I and II, leading to receptor cross-linking and signal transduction. The receptors differ strongly in their intra-cellular signaling pathways. TNF-alpha was originally described as a highly cytotoxic cytokine for tumor cells, it causes tumor necrosis in vivo and shows cytolytic activity against tumor cells in vitro. Furthermore, TNF-alpha is found to be a central mediator in many inflammatory and immunological processes. It can be induced by various products of micro-organisms and by various cytokines leading to expression of a wide variety of cytokines. The pro-inflammatory properties of TNF-alpha play a central role in several auto-immune diseases such as rheumatoid arthritis and inhibition by neutralizing molecules have been shown to be beneficial in patients.

References

1. Bradding; P et al. Am J Respir Cell Mol Biol 1994; 10: 471
2. Bradding, P et al: Clin Exp Allergy 1995, 25: 406
3. Gerspach; J et al. Microsc Res Tech 2000; 50: 243
4. Laan van der N et al. Arch Dermatol Res 2001; 293: 226
5. Lee E et al. Ann Dermatol 2009; 21:345

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES