



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-IFN-Alpha (II), clone N27 (Monoclonal)

Clone no. N27

MONOSAN

Product name	Mouse anti-IFN-Alpha (II), clone N27 (Monoclonal)
Host	Mouse
Applications	ELISA, IHC-fr, WB
Species reactivity	human
Conjugate	-
Immunogen	E.coli derived recombinant human IFNalpha2c
Isotype	IgG1,kappa
Clonality	Monoclonal
Clone number	N27
Size	100 ug
Concentration	100 ug/ml
Format	-
Storage buffer	PBS with 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-IFN-Alpha (II), clone N27 (Monoclonal)

Clone no. N27

MONOSAN

Additional info

The alpha interferons are involved in virus resistance in target cells for these viruses. They are known to block cell proliferation and to regulate MHC class I antigen expression. The IFN α family has over 20 genes and pseudogenes in two families (I and II), one with a mature length of 166aa and one of 172aa. Cells producing IFN α are lymphocytes, monocytes, macrophages and cell lines such as Namalwa and KGI. Bioassays for IFN α include cytopathic effect blocking, by viruses such as VSV, SFV and BMCV, on their target cells. A number of receptors for IFN α are now known and seem to be expressed on most cell types. N27 is specific for human IFN α 2 and does not cross react with human IFN α 1. N27 reacts with linear peptide 43aa-53aa, placing the epitope outside the immunodominant regions I and II.

References

1. Kontsek, P. et al., Mol Immunol. 29: 863-870 (1992)
2. Kontsek, P. et al., Immunol. Lett. 35: 281-284 (1993)
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES