



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Mouse anti-VCAM-1, clone 1G11B1 (Monoclonal)

Clone no. 1G11B1

MONOSAN

Product name Mouse anti-VCAM-1, clone 1G11B1 (Monoclonal)

Host Mouse

Applications IHC-fr,FC,FUNC,IF,IP

Species reactivity human

Conjugate -

Immunogen Unknown or proprietary to MONOSAN and/or its suppliers

Isotype IgG1

Clonality Monoclonal

Clone number 1G11B1

Size 1 ml

Concentration 100 ug/ ml

Format -

Storage buffer PBS with 0.1% BSA and 0.02% sodium azide

Storage until expiry date 2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-VCAM-1, clone 1G11B1 (Monoclonal)

Clone no. 1G11B1

MONOSAN

Additional info

The monoclonal antibody 1G11B1 recognizes vascular cell adhesion molecule-1 (VCAM-1). VCAM-1 is a member of the immunoglobulin superfamily of adhesion molecules, which includes ICAMs, PECAM-s and MADCAM, and is involved in leukocyte-endothelial cell interactions. The immunoglobulin superfamily is a type I transmembrane protein characterized by extracellular immunoglobulin domains, a transmembrane region and a cytoplasmic tail. They are essential for the development of the embryo and for immune and inflammatory responses. These transmembrane glycoproteins mediate cell interaction with, and adhesion to, other cells and the extracellular matrix. VCAM-1 contains six immunoglobulin domains of the H-type and interacts with VLA-4 expressed on leukocytes. Multiple adhesion molecules play a role in leukocyte recruitment. The process of migration of a leukocyte through the vascular endothelium consists of the following steps: leukocyte-endothelium interaction (first tethering and rolling and then adhesion) and transendothelial migration. VCAM-1 is almost not expressed under physiological conditions. However, under appropriate pro-inflammatory conditions where the endothelium is exposed to inflammatory cytokines such as tumour necrosis factor- α or IL-1 β and becomes activated, VCAM-1 gene expression is rapidly elevated by the vascular endothelium. There is also a soluble form of VCAM-1 which is angiogenic and chemotactic for endothelial cells. sVCAM-1 is up-regulated in several disease states (eg, myocardial infarction, type 2 diabetes mellitus, primary antiphospholipid syndrome, and rheumatoid arthritis).

References

1. Thornhill; M et al. J Immunology 1991; 146: 592
2. Van der Vieren, M et al. J Immunol 1999, 163:1984
3. Dienst; A et al. J Natl Cancer Inst 2005; 97 :733
4. Wijbrandt C et al. Ann Rheum Dis 2008; 67:1139
5. Roussel L et al. J Immunol 210; 184; 4531

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES