



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Clusterin, clone Hs-3 (Monoclonal)

Clone no. Hs-3

MONOSAN

Product name	Mouse anti-Clusterin, clone Hs-3 (Monoclonal)
Host	Mouse
Applications	WB, IHC-P, ICC, ELISA
Species reactivity	Human
Conjugate	-
Immunogen	Freshly ejaculated human sperms were washed in PBS and extracted in 3% acetic acid, 10% glycerol, 30 mM benzaminidine. The acid extract was
Isotype	IgG1
Clonality	Monoclonal
Clone number	Hs-3
Size	0.1 mg
Concentration	1 mg/ml
Format	-
Storage buffer	Phosphate buffered saline (PBS) solution with 15 mM sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Clusterin, clone Hs-3 (Monoclonal)

Clone no. Hs-3

MONOSAN

Additional info

Clusterin (APO J, SGP-2, TRPM-2, SP-40, pADHC-9, CLJ, T64, GP III, XIP8) is a 75-80 kD disulfide-linked heterodimeric protein containing about 30% of N-linked carbohydrate rich in sialic acid but truncated forms targeted to the nucleus have also been identified. It is a conserved secreted glycoprotein expressed by a wide range of tissues and being implicated in many physiological processes, including e.g. lipid transportation, complement inhibition, tissue remodeling, membrane recycling, or clearance of cellular debris. It is nearly ubiquitously expressed in most mammalian tissues and can be found in plasma, milk, urine, cerebrospinal fluid and semen. Clusterin is able to bind and form complexes with numerous partners (immunoglobulins, lipids, heparin, bacteria, complement components, paraoxonase, beta amyloid, leptin etc.) and is expressed in many pathological and clinically relevant situations including cancer, organ regeneration, infection, Alzheimer disease, retinitis pigmentosa, myocardial infarction, renal tubular damage, autoimmunity and others. A genuine function of clusterin is still enigmatic.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES