



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rabbit anti-TCL1, clone EP105 (Monoclonal)

Clone no. EP105

MONOSAN

Product name	Rabbit anti-TCL1, clone EP105 (Monoclonal)
Host	Rabbit
Applications	IHC-P (1:100-1:200)
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG
Clonality	Monoclonal
Clone number	EP105
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	Tris Buffer, pH 7.3-7.7, containing 1% BSA and <0.1% Sodium Azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Rabbit anti-TCL1, clone EP105 (Monoclonal)

Clone no. EP105

MONOSAN

Additional info

T-cell leukemia/lymphoma protein 1A (TCL1) is a member of theTCL1 family and enhances the phosphorylation and activation ofAKT1, AKT2 and AKT3. TCL1promotes the nuclear translocation ofAKT1 and enhances cell proliferation, stabilizes mitochondrial membrane potential and promotes cell survival. The expression of TCL1 is restricted to lymphoid cells. It is expressed early in lymphocyte differentiation. Strong expression ofTCL1 is found in a subset of mantle zone B lymphocytes and is expressed to a lesser extent by follicle center cells. In B cell neoplasia, TCL1 immunoreactivity is found in the majority of B cell lymphomas including lymphoblastic lymphoma, chronic lymphocytic leukemia, mantle cell lymphoma, follicular lymphoma, Burkitt lymphoma, diffuse large B-cell lymphoma (60%), and primary cutaneous B cell lymphoma (55%). The expression of theTCL1 genecharacterizes low-grade B cell lymphomas.

References

1. Laine J, et al. Mol Cell2000, 6:395-407
2. Pekarsky Y, et al. Proc Natl Acad Sci U S A, 2000, 97:3028-3033
3. Narducci MG, et al. Cancer Res, 2000, 60:2095-2100
4. Roos J, et al. Pathobiology, 2001, 69:59-66
5. Nakayama I, et al. Pathol Int, 2000, 50:191-199

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES