



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Lactoferrin, clone 265-1K1 (Monoclonal)

Clone no. 265-1K1

MONOSAN

Product name	Mouse anti-Lactoferrin, clone 265-1K1 (Monoclonal)
Host	Mouse
Applications	IHC-fr,ELISA,WB
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	265-1K1
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Lactoferrin, clone 265-1K1 (Monoclonal)

Clone no. 265-1K1

MONOSAN

Additional info

The monoclonal antibody 265-1K1 reacts with human lactoferrin (LF), an 80 kDa glycoprotein. Lactoferrin was first isolated from human milk and plays an important part in the immune system and helps to fight infections. Lactoferrin promotes the health of the gastro-intestinal system by improving the intestinal microbial balance. In addition, LF can be found in epithelia and most body fluids and secretions. Lactoferrin is secreted in plasma by neutrophils. Its plasma concentration also represents a positive relation to the total pool of neutrophils and the rate of neutrophil turnover. In inflammation lactoferrin is released from secondary granules of neutrophilic leukocytes into the extracellular medium. Therefore the extracellular lactoferrin concentration can be used as an index for neutrophil activation. Lactoferrin strongly binds to iron and this iron binding property is considered to be an important antimicrobial. Human lactoferrin binds to bacterial products through its highly positively charged N-terminus, it kills various bacteria, most probably by inducing intracellular changes in these bacteria without affecting the membrane permeability. Cleavage by pepsin of lactoferrin leads to the release of lactoferricin H. This 47 amino acid peptide has more antimicrobial activity than its precursor and it can inhibit the classical but not the alternative complement pathway. Lactoferrin also plays a role in signal transduction, immunomodulation and has antiadhesive, anticancer, antiviral activity.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES