



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Mouse anti-Elastase, clone 265-3K1 (Monoclonal)

Clone no. 265-1K1

MONOSAN

Product name	Mouse anti-Elastase, clone 265-3K1 (Monoclonal)
Host	Mouse
Applications	FC,ELISA,IF,IHC-P,WB
Species reactivity	human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	265-1K1
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Elastase, clone 265-3K1 (Monoclonal)

Clone no. 265-1K1

MONOSAN

Additional info

The monoclonal antibody 265-3K1 recognizes human leukocyte elastase. Leukocyte elastase, a major serine proteinase in man, is predominantly present in the azurophilic granules of neutrophils and monocytes. Elastase has a broad range of extracellular matrix substrates including elastin, proteoglycans, collagen and fibronectin. The action of elastase is controlled by serine proteinase inhibitors. Elastase, when released during inflammation, is rapidly bound by its two main inhibitors, alpha1-PI and alpha2-macroglobuline to form elastase-inhibitor complexes. In addition mucosa secretions may contain the locally secreted elastase inhibitors elafin/SKALP and SLPI. When secreted at sites of inflammation elastase can cause severe tissue damage. An important role has been suggested for human elastase in various inflammatory disorders, including pulmonary emphysema, sepsis, arthritis, nephritis and certain skin diseases. Elastase induces the production of IL-8 in human bronchial epithelial, a proces that occurs in part through TLR4.

References

1. Davies; P et al. J Immunol Meth 2008; 366:175
2. Delbosc, S et al PLos ONE 2011, 6: e18679
3. Novak; T et al. Thesis 2013
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES