



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Rat ASGPR, clone 8D7 (Monoclonal)

Clone no. 8D7

MONOSAN

Product name	Mouse anti-Rat ASGPR, clone 8D7 (Monoclonal)
Host	Mouse
Applications	IHC-fr,FC,ELISA,IF,WB
Species reactivity	rat, human
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	8D7
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Rat ASGPR, clone 8D7 (Monoclonal)

Clone no. 8D7

MONOSAN

Additional info

The asialoglycoprotein (ASGP) receptor is a transmembrane hepatocellular surface carbohydrate binding glycoproteins lacking terminal sialic acid residues (asialoglycoproteins). Characterization of the ASGP receptor-revealed its functional role in the binding, internalization and transport of a wide range of glycoproteins, which have exposed galactose or N-acetylgalactosamine residues, via the process of receptor-mediated endocytosis (RME). The ASGP receptor can bind a variety of important plasma proteins including transport proteins (i.e. transferrin), enzymes such as alkaline phosphatase, immunoglobulins including IgA, apoptotic hepatocytes, fibronectin and platelets. Additionally, the expression of the ASGP receptor has been clinically correlated to the level of hepatic function that is lost during liver diseases related to cancer, viral hepatitis, and cirrhosis. The ASGP receptor consists of major and minor subunits, which in the rat were identified as rat hepatic lectin (RHL) 1 and RHL 2/3, with molecular weights of respectively 42, 49 and 54 kDa. The selective binding (calcium and pH depended) and uptake of terminal galactosyl bearing proteins requires the formation of hetero-oligomers between these major and minor forms. The total ASGP receptor population consisted of two functionally distinct receptor populations, designated State 1 and State 2, which were involved in the endocytosis and intracellular processing of ligands by different pathways. The monoclonal antibody 8D7 recognizes a subunit-specific epitope on RHL-1 of rat ASGPR. The monoclonal antibody 8D7 is cross reactive with human ASGPR.

References

1. Mizuno; M et al. Gastroenterol Japan 1986; 21: 238
2. Mizuno, M et al Liver 1987, 7: 251
3. Hyodo; I et al. Liver 1993; 13: 80
4. Shimada M et al. Hepatol Res 2003; 26: 55
5. HiraiM et al. Acta med okayama 2002 56: 135

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES