



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Shigatoxin 1, clone 13C4 (Monoclonal)

Clone no. 13C4

MONOSAN

---

Product name	Mouse anti-Shigatoxin 1, clone 13C4 (Monoclonal)
Host	Mouse
Applications	FUNC,ELISA,WB
Species reactivity	n/a
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	13C4
Size	1 ml
Concentration	100 ug/ ml
Format	-
Storage buffer	PBS with 0.1% BSA and 0.02% sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti-Shigatoxin 1, clone 13C4 (Monoclonal)

Clone no. 13C4

MONOSAN

**Additional info**

The monoclonal antibody 13C4 recognizes the 1B subunit of Shiga-like toxin 1. Shiga-like toxins (SLTs), are also called Verotoxins. Enterohemorrhagic Escherichia coli (EHEC) strains which are primarily of serotypes O157:H7, O26:H11 and O111:H8 have been incriminated as etiologic agents of hemorrhagic colitis and Hemolytic-uremic syndrome, a generalized disease characterized by acute renal failure, thrombocytopenia, and microangiopathic hemolytic anemia. There are several distinct E.coli SLTs. SLT-I and SLT-II are produced by EHEC. SLT-I and Shiga toxin share 99% deduced amino acid sequence homology, whereas SLT-I and SLT-II share about 60% deduced amino acid sequence homology. SLT-I and SLT-II are antigenically distinct. The protein structure of the toxin consists of two domains: the A polypeptide that inhibits protein synthesis by targeting ribosomes, and the B polypeptide pentamer that binds to the eukaryotic cell receptor globotriaosylceramide (Gb3) leading to receptor-mediated endocytosis.

**References**

1. Strockbine; N et al. Infect Immun 1985; 50: 695
2. Smith, M et al Vaccine 2006, 24: 4122
3. Smith; M et al. Infect Immun 2006; 74: 6992
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES