



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-VCP (Intra-acrosomal protein), clone Hs-14 (Monoclonal)

Clone no. Hs-14

MONOSAN

Product name	Mouse anti-VCP (Intra-acrosomal protein), clone Hs-14 (Monoclonal)
Host	Mouse
Applications	FC , WB, ICC
Species reactivity	Mouse, Human
Conjugate	-
Immunogen	Freshly ejaculated human sperms were washed in PBS and extracted in 3% acetic acid, 10% glycerol, 30 mM benzaminidine. The acid extract was
Isotype	IgM
Clonality	Monoclonal
Clone number	Hs-14
Size	0.1 mg
Concentration	1 mg/ml
Format	-
Storage buffer	Tris buffered saline (TBS) solution with 15 mM sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-VCP (Intra-acrosomal protein), clone Hs-14 (Monoclonal)

Clone no. Hs-14

MONOSAN

Additional info

VCP (valosin-containing protein), also known as p97, TERA, ALS14, IBMPFD, HEL-220, IBMPFD1, or HEL-S-70, is a member of a protein family that includes putative ATP-binding proteins involved in vesicle transport and fusion, 26S proteasome function, and assembly of peroxisomes. VCP is a structural protein that associates with clathrin and heat-shock protein Hsc70, to form a complex. It has been implicated in a number of cellular events that are regulated during mitosis, including homotypic membrane fusion, spindle pole body function, and ubiquitin-dependent protein degradation. In sperm this intra-acrosomal protein can be used as a marker for evaluation of the physiological state of sperm cells as well as for selection of a suitable method of fertilization in the laboratories of assisted reproduction.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES