



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Product datasheet MON8070

MONOSAN[®]

Mouse anti-CD55, clone 67 (Monoclonal)

Clone no. 67

MONOSAN

Product name Mouse anti-CD55, clone 67 (Monoclonal)

Host Mouse

Applications WB,FC,IHC-fr

Species reactivity human

Conjugate Purified

Immunogen K562 cells

Isotype IgG1

Clonality Monoclonal

Clone number 67

Size 0.1 mg

Concentration 1.0 mg/ml

Format -

Storage buffer PBS with 0.09% sodium azide

Storage until expiry date 2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-CD55, clone 67 (Monoclonal)

Clone no. 67

MONOSAN

Additional info

Mouse anti Human CD55 antibody, clone 67 recognizes the human CD55 cell surface antigen, a GPI linked molecule also known as decay accelerating factor (DAF). CD55 is expressed by a wide range of cell types. CD55 is the complement regulatory protein, decay accelerating factor (DAF) (Lublin and Atkinson 1989). Human CD55 is a 126;70 kDa glycoprotein (in erythrocytes) anchored in the membrane by glycosylphosphatidylinositol tail. In other cells the apparent molecular weight is somewhat larger. It has a substantial content of O-glycans, and also on N-glycan. DAF binds to activated C4b or C3b complement fragments on the cell surface, preventing the assembly and accelerating the decay of both classical and alternative pathways. DAF carries the Cromer related blood group antigens. DAF has a wide distribution on cells in non-haematopoietic tissues, particularly epithelium and is found at the fetal-maternal interface in placenta (Holmes et al. 1990 and Yang et al. 2009). Soluble forms of DAF are found, for example, in plasma, saliva and urine (Medof et al. 1987). The antigen on erythrocytes is pronase and chymotrypsin sensitive, but resistant to trypsin.

References

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES