



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Mouse anti-Neural Cell Adhesion Molecule, NCAM, clone RNL-1 (Monoclonal)

Clone no. RNL-1

MONOSAN

---

|                           |  |
|---------------------------|--|
| Product name              | Mouse anti-Neural Cell Adhesion Molecule, NCAM, clone RNL-1 (Monoclonal) |
| Host                      | Mouse  |
| Applications              | IHC-fr (1:10-1:20), WB   |
| Species reactivity        | Human  |
| Conjugate                 | -  |
| Immunogen                 | Cytoskeletal preparation of human epidermis                              |
| Isotype                   | IgG1   |
| Clonality                 | Monoclonal   |
| Clone number              | RNL-1  |
| Size                      | 1 ml   |
| Concentration             | n/a  |
| Format                    | -  |
| Storage buffer            | Contains 0.09% sodium azide  |
| Storage until expiry date | 2-8°C  |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

Mouse anti-Neural Cell Adhesion Molecule, NCAM, clone RNL-1 (Monoclonal)

Clone no. RNL-1

MONOSAN

---

**Additional info**

Mab RNL-1, directed against N-CAM (Neural Cell Adhesion Molecule), reacts with normal neural tissues and endocrine glands, such as pancreatic islet, pituitary gland and adrenal medulla. Expression was also found in Leydig cells of the testis, in the thyroid and in smooth-muscle cells of the small intestine, colon and bladder. The antibody is a valuable marker for the characterization of several neuroendocrine tumors. In immunoblotting (Western) the antibody is reactive with 3 main clusters of protein bands in the molecular weight region of 200 kD, 100 kD, and 25-27 kD. Positive control: Pancreatic islet cells (cell surface staining).

|            |    |   |
|------------|----|---|
| References | 1. | - |
|            | 2. | - |
|            | 3. | - |
|            | 4. | - |
|            | 5. | - |

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES